

Title	Relating Bishop's function spaces to neighbourhood spaces
Author(s)	Ishihara, Hajime
Citation	Annals of Pure and Applied Logic, 164(4): 482-490
Issue Date	2012-11-06
Type	Journal Article
Text version	author
URL	http://hdl.handle.net/10119/11406
Rights	NOTICE: This is the author's version of a work accepted for publication by Elsevier. Hajime Ishihara, Annals of Pure and Applied Logic, 164(4), 2012, 482-490, http://dx.doi.org/10.1016/j.apal.2012.10.009
Description	

Relating Bishop's function spaces to neighbourhood spaces

Hajime Ishihara

April 24, 2011

Abstract

We extend Bishop's concept of function spaces to the concept of pre-function spaces. We show that there is an adjunction between the category of neighbourhood spaces and the category of Φ -closed pre-function spaces. We also show that there is an adjunction between the category of uniform spaces and the category of Ψ -closed pre-function spaces.

Keywords: constructive mathematics, function space, neighbourhood space, uniform space, completeness, cocompleteness, adjunction.

2000 Mathematics Subject Classification: 03F65, 54E05.

1 Introduction

In 1967, Bishop [3] proposed two approaches to topology in his constructive mathematics: one approach is based on the idea of a neighbourhood space, and the other is based on the idea of a function space. However, in his book, he did not investigate them in detail.

It turns out that neighbourhood spaces are both formal topologies, as introduced by Sambin [19, 20, 21], and constructive topological spaces (see Aczel [1]). In addition, connections between neighbourhood spaces and other constructive topological notions – in particular the Bridges-Vîță one of an apartness space [7, 9] – have been explored [14, 13]. On the other hand, the approach to constructive topology based on the idea of a function space has lain relatively dormant for over forty years.

Recently, Bridges [5] has dealt with various aspects of function spaces which revive Bishop's approach to topology based on function spaces. Following Bishop [3, Definition 8, Chapter 3], we define a *function space* X to be a pair $(\underline{X}, \mathcal{F}_X)$ of a set \underline{X} and a set \mathcal{F}_X of functions from \underline{X} to \mathbf{R} satisfying the following conditions.

- F1. \mathcal{F}_X contains the constant functions.
- F2. Sums and products of elements of \mathcal{F}_X are in \mathcal{F}_X .
- F3. The composition $\varphi \circ f$ of an element f of \mathcal{F}_X and a continuous function $\varphi : \mathbf{R} \rightarrow \mathbf{R}$ is in \mathcal{F}_X , where $\varphi : \mathbf{R} \rightarrow \mathbf{R}$ is continuous if it is uniformly continuous on every compact interval.
- F4. Uniform limits of elements of \mathcal{F}_X are in \mathcal{F}_X ; that is, if for each $\epsilon > 0$ there exists g in \mathcal{F}_X , with $|g(x) - f(x)| \leq \epsilon$ for all x in \underline{X} , then $f \in \mathcal{F}_X$.

Bishop called \mathcal{F}_X the *topology* on \underline{X} .

In this paper, we first introduce the notion of a pre-function space just as a pair of a set S and a set of real-valued functions on S , and the notion of a function space morphism according to [5]. Then we focus on the condition F3 above, and introduce the notion of a C -complete pre-function space for a set C of functions from \mathbf{R} to \mathbf{R} ; in the definition of a function space, C is taken to be the set of continuous functions in the above sense. We show that the category of C -complete pre-function spaces with function space morphisms is complete and cocomplete.

We propose a closure condition Φ_S on a set of real-valued functions on a set S , and introduce the notion of a Φ -closed pre-function space. It emerges that each Φ -closed pre-function space is a function space in Bishop's sense. Then we construct an adjunction between the category of neighbourhood spaces with continuous functions in usual sense and the category of Φ -closed pre-function spaces with function space morphisms, which relates Bishop's two approaches to topology, and show that the category of Φ -closed pre-function spaces is complete and cocomplete. We also construct an adjoint equivalence between the category of neighbourhood spaces with a compatible family of pseudometrics and the category of Φ -closed pre-function spaces.

Finally, we introduce another closure condition Ψ_S on a set of real-valued functions on a set S and the corresponding notion of a Ψ -closed pre-function space, and construct an adjunction between the category of

uniform spaces with uniformly continuous functions and the category of Ψ -closed pre-function spaces with function space morphisms.

Although the results are presented in informal Bishop-style constructive mathematics [3, 4, 6, 22, 8], it is possible to formalize them in Aczel's constructive Zermelo-Fraenkel set theory (**CZF**) [2] together with Relativized Dependent Choice (RDC).

There are other constructive treatments of topology: see, for example, Grayson [11, 12].

2 Complete pre-function spaces

A *pre-function space* X is a pair $(\underline{X}, \mathcal{F}_X)$ consisting of a set \underline{X} and a set \mathcal{F}_X of functions from \underline{X} to \mathbf{R} , called a *function space structure* on \underline{X} . According to [5], a *function space morphism* from a pre-function space X into a pre-function space Y is a mapping $f : \underline{X} \rightarrow \underline{Y}$ such that

$$\forall g \in \mathcal{F}_Y (g \circ f \in \mathcal{F}_X).$$

We write $f : X \rightarrow Y$ to denote that f is a function space morphism from X into Y , and $\text{Hom}(X, Y)$ for the set of function space morphisms from X into Y .

For any set S , there are the pre-function spaces (S, \mathbf{R}^S) , where \mathbf{R}^S is the set of functions from S into \mathbf{R} , and (S, \emptyset) , called the *discrete* function space of S and the *trivial* pre-function space of S , respectively. For each pre-function space Y , any mapping $f : S \rightarrow \underline{Y}$ is a function space morphism from the discrete function space of S into Y , and any mapping $f : \underline{Y} \rightarrow S$ is a function space morphism from Y into the trivial pre-function space of S .

Let C be a set of functions from \mathbf{R} to \mathbf{R} containing the identity map $\text{id}_{\mathbf{R}}$ and closed under composition. A pre-function space X is *C-complete* if

$$\forall f \in \mathcal{F}_X \forall \varphi \in C (\varphi \circ f \in \mathcal{F}_X).$$

The discrete function spaces and the trivial pre-function spaces are C -complete for any C , and any pre-function space is $\{\text{id}_{\mathbf{R}}\}$ -complete. If a pre-function space X is C -complete, then X is C' -complete for any $C' \subseteq C$. Since C is closed under composition, the pre-function space $\mathbf{R}_C = (\mathbf{R}, C)$ is C -complete.

Lemma 2.1. *Let X be a pre-function space. Then*

1. $\text{Hom}(X, \mathbf{R}_C) \subseteq \mathcal{F}_X$,
2. X is C -complete if and only if $\mathcal{F}_X \subseteq \text{Hom}(X, \mathbf{R}_C)$,
3. X is C -complete if and only if $\mathcal{F}_X = \text{Hom}(X, \mathbf{R}_C)$.

Proof. Straightforward. For (1), note that $\text{id}_{\mathbf{R}} \in C$. □

Especially, $C = \text{Hom}(\mathbf{R}_C, \mathbf{R}_C)$.

Proposition 2.2. *Let X be a pre-function space. Then the pre-function space $\tilde{X} = (\underline{X}, \text{Hom}(X, \mathbf{R}_C))$, called the C -completion of X , is C -complete. Furthermore, $\text{id}_{\underline{X}} : X \rightarrow \tilde{X}$, and if Y is a C -complete pre-function space and $f : X \rightarrow Y$, then $f : \tilde{X} \rightarrow Y$.*

Proof. Let $\varphi \in C$ and $f \in \text{Hom}(X, \mathbf{R}_C)$. Then for each $\psi \in C$, since $\psi \circ (\varphi \circ f) = (\psi \circ \varphi) \circ f$ and $\psi \circ \varphi \in C$, we have $\psi \circ (\varphi \circ f) \in \mathcal{F}_X$, and therefore $\varphi \circ f \in \text{Hom}(X, \mathbf{R}_C)$. Hence \tilde{X} is C -complete.

Since $\text{Hom}(X, \mathbf{R}_C) \subseteq \mathcal{F}_X$, by Lemma 2.1 (1), we have $\text{id}_{\underline{X}} : X \rightarrow \tilde{X}$. Let Y be a C -complete pre-function space, and let $f : X \rightarrow Y$. Then for each $g \in \mathcal{F}_Y$ and $\varphi \in C$, since $\varphi \circ g \in \mathcal{F}_Y$, we have $\varphi \circ (g \circ f) = (\varphi \circ g) \circ f \in \mathcal{F}_X$, and therefore $g \circ f \in \text{Hom}(X, \mathbf{R}_C)$. Hence $f : \tilde{X} \rightarrow Y$. □

Let \mathbf{Func}_C denote the category of function spaces whose objects are C -complete pre-function spaces and whose morphisms are function space morphisms. For basic notions and results in category theory, we refer the reader to [10, 16, 17, 18].

Note that, in \mathbf{Func}_C , the initial object is the discrete function space of \emptyset , the terminal objects are the trivial pre-function spaces of singletons.

Let \mathbf{I} and \mathbf{C} be categories. A *cone* of a functor $H : \mathbf{I} \rightarrow \mathbf{C}$ is an object L in \mathbf{C} , together with a family of morphisms $\phi_I : L \rightarrow H(I)$ for each object I in \mathbf{I} , such that $H(i) \circ \phi_I = \phi_J$ for each morphism $i : I \rightarrow J$ in \mathbf{I} . A cone $\langle L, \phi_I \rangle$ of a functor $H : \mathbf{I} \rightarrow \mathbf{C}$ is a *limit* of H if for each cone $\langle X, \psi_I \rangle$ of H there exists a unique morphism $u : X \rightarrow L$ such that $\phi_I \circ u = \psi_I$ for each object I in \mathbf{I} . We say that \mathbf{C} is *complete* if every functor $H : \mathbf{I} \rightarrow \mathbf{C}$ from a small category \mathbf{I} has a limit. A *cocone* of a functor $H : \mathbf{I} \rightarrow \mathbf{C}$ is an object L in \mathbf{C} , together with a family of morphisms $\phi_I : H(I) \rightarrow L$ for each object I in \mathbf{I} , such that $\phi_J \circ H(i) = \phi_I$ for each morphism $i : I \rightarrow J$ in \mathbf{I} . A cocone $\langle L, \phi_I \rangle$ of a functor $H : \mathbf{I} \rightarrow \mathbf{C}$ is a *colimit* of H if for each cocone $\langle X, \psi_I \rangle$ of H there exists a unique morphism $u : L \rightarrow X$ such that $u \circ \phi_I = \psi_I$ for each

object I in \mathbf{I} . We say that \mathbf{C} is *cocomplete* if every functor $H : \mathbf{I} \rightarrow \mathbf{C}$ from a small category \mathbf{I} has a colimit.

We will show that the category \mathbf{Fun}_C is complete and cocomplete.

Proposition 2.3. *Let S be a set, let $\{X_i\}_{i \in I}$ be a family of C -complete pre-function spaces, and for each $i \in I$ let $f_i : S \rightarrow X_i$. Then there exists a function space structure \mathcal{F} on S such that the pre-function space (S, \mathcal{F}) is C -complete, and if h is a mapping from the underlying set \underline{Y} of a pre-function space Y into S , then $h : Y \rightarrow (S, \mathcal{F})$ if and only if $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$.*

Proof. Let

$$\mathcal{F} = \{f \circ f_i \mid i \in I, f \in \mathcal{F}_{X_i}\}.$$

Then for each $\varphi \in C$, $i \in I$ and $f \in \mathcal{F}_{X_i}$, since $\varphi \circ (f \circ f_i) = (\varphi \circ f) \circ f_i$ and $\varphi \circ f \in \mathcal{F}_{X_i}$, we have $\varphi \circ (f \circ f_i) \in \mathcal{F}$, and hence the pre-function space (S, \mathcal{F}) is C -complete.

It is straightforward to see that $f_i : (S, \mathcal{F}) \rightarrow X_i$ for each $i \in I$, and hence if $h : Y \rightarrow (S, \mathcal{F})$, then $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. Let Y be a pre-function space and let $h : \underline{Y} \rightarrow S$ be such that $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. Then for each $i \in I$ and $f \in \mathcal{F}_{X_i}$, we have $(f \circ f_i) \circ h = f \circ (f_i \circ h) \in \mathcal{F}_Y$, and hence $h : Y \rightarrow (S, \mathcal{F})$. \square

Proposition 2.4. *Let S be a set, let $\{X_i\}_{i \in I}$ be a family of C -complete pre-function spaces, and for each $i \in I$ let $f_i : X_i \rightarrow S$. Then there exists a function space structure \mathcal{F} on S such that the pre-function space (S, \mathcal{F}) is C -complete, and if h is a mapping from S into the underlying set \underline{Y} of a pre-function space Y , then $h : (S, \mathcal{F}) \rightarrow Y$ if and only if $h \circ f_i : X_i \rightarrow Y$ for each $i \in I$.*

Proof. Let

$$\mathcal{F} = \{f \in \mathbf{R}^S \mid \forall i \in I (f \circ f_i \in \mathcal{F}_{X_i})\}.$$

Then for each $f \in \mathcal{F}$ and $\varphi \in C$, since $(\varphi \circ f) \circ f_i = \varphi \circ (f \circ f_i) \in \mathcal{F}_{X_i}$, we have $\varphi \circ f \in \mathcal{F}$, and hence the pre-function space (S, \mathcal{F}) is C -complete.

It is straightforward to see that $f_i : X_i \rightarrow (S, \mathcal{F})$ for each $i \in I$, and hence if $h : (S, \mathcal{F}) \rightarrow Y$, then $h \circ f_i : X_i \rightarrow Y$ for each $i \in I$. Let Y be a pre-function space and let $h : S \rightarrow \underline{Y}$ be such that $h \circ f_i : X_i \rightarrow Y$ for each $i \in I$. Then for each $g \in \mathcal{F}_Y$, since $(g \circ h) \circ f_i = g \circ (h \circ f_i) \in \mathcal{F}_{X_i}$ for each $i \in I$, we have $g \circ h \in \mathcal{F}$, and hence $h : (S, \mathcal{F}) \rightarrow Y$. \square

Theorem 2.5. *The category \mathbf{Fun}_C is complete and cocomplete.*

Proof. Let $H : \mathbf{I} \rightarrow \mathbf{Fun}_C$ be a functor from a small category \mathbf{I} . Then, since the category \mathbf{Set} of sets and mappings is complete, the functor $KH : \mathbf{I} \rightarrow \mathbf{Set}$ has a limit $\langle S, \phi_I \rangle$ in \mathbf{Set} , where K is the forgetful functor from \mathbf{Fun}_C into \mathbf{Set} , taking each C -complete pre-function space X to its underlying set \underline{X} and each function space morphism to itself. By Proposition 2.3, since $\phi_I : S \rightarrow \underline{H(I)}$ for each object I in \mathbf{I} , there exists a function space structure \mathcal{F} on S such that $L = (S, \mathcal{F})$ is an object in \mathbf{Fun}_C , and if h is a mapping from the underlying set \underline{Y} of an object Y in \mathbf{Fun}_C into S , then $h : Y \rightarrow L$ if and only if $\phi_I \circ h : Y \rightarrow H(I)$ for each object I in \mathbf{I} .

Since $\text{id}_S : L \rightarrow L$, we have $\phi_I = \phi_I \circ \text{id}_S : L \rightarrow H(I)$ for each object I in \mathbf{I} , and hence $\langle L, \phi_I \rangle$ is a cone of H in \mathbf{Fun}_C . Let $\langle Y, \psi_I \rangle$ be a cone of H in \mathbf{Fun}_C . Then $\langle \underline{Y}, \psi_I \rangle$ is a cone of KH in \mathbf{Set} , and hence there exists a unique mapping $h : \underline{Y} \rightarrow S$ such that $\phi_I \circ h = \psi_I$ for each object I in \mathbf{I} . Therefore, since $\phi_I \circ h = \psi_I : Y \rightarrow H(I)$ for each object I in \mathbf{I} , we have $h : Y \rightarrow L$. Thus $\langle L, \phi_I \rangle$ is a limit of H in \mathbf{Fun}_C .

Similarly, using Proposition 2.4 instead of Proposition 2.3, we see that \mathbf{Fun}_C is cocomplete. \square

3 Neighbourhood and function spaces

A *neighbourhood space* A is a pair (\underline{A}, τ_A) consisting of a set \underline{A} and an inhabited set τ_A of subsets of \underline{A} , called an *open base* on \underline{A} , such that

$$\text{NS1. } \forall x \in \underline{A} \exists U \in \tau_A (x \in U),$$

$$\text{NS2. } \forall x \in \underline{A} \forall U, V \in \tau_A [x \in U \cap V \implies \exists W \in \tau_A (x \in W \subseteq U \cap V)];$$

see [3, Chapter 3]. A *continuous mapping* f from a neighbourhood space A into a neighbourhood space B is a mapping $f : \underline{A} \rightarrow \underline{B}$ such that

$$\forall x \in \underline{A} \forall V \in \tau_B [f(x) \in V \implies \exists U \in \tau_A (x \in U \subseteq f^{-1}(V))].$$

We write $f : A \rightarrow B$ to denote that f is a continuous mapping from A into B , and $\mathcal{C}(A, B)$ for the set of continuous mappings from A into B .

For any set S , there are the neighbourhood spaces (S, σ_S) , where σ_S is the set of singletons of S , and $(S, \{S\})$, called the *discrete* neighbourhood space of S and the *trivial* neighbourhood space of S , respectively. For each

neighbourhood space Y , any mapping $f : S \rightarrow \underline{Y}$ is a continuous mapping from the discrete neighbourhood space of S into Y , and any mapping $f : \underline{Y} \rightarrow S$ is a continuous mapping from Y into the trivial neighbourhood space of S .

Let \mathbf{Nbh} denote the category of neighbourhood spaces whose objects are neighbourhood spaces and whose morphisms are continuous mappings. Note that, in \mathbf{Nbh} , the initial object is the discrete neighbourhood space of \emptyset , the terminal objects are the trivial neighbourhood spaces of singletons.

As we will see in Theorem 3.3, the category \mathbf{Nbh} is complete and cocomplete.

Proposition 3.1. *Let S be a set, let $\{A_i\}_{i \in I}$ be a family of neighbourhood spaces, and for each $i \in I$ let $f_i : S \rightarrow \underline{A_i}$. Then there exists an open base τ on S such that if h is a mapping from the underlying set \underline{B} of a neighbourhood space B into S , then $h : B \rightarrow (S, \tau)$ if and only if $f_i \circ h : B \rightarrow A_i$ for each $i \in I$.*

Proof. Let

$$\tau = \{\bigcap_{k=1}^n f_{i_k}^{-1}(U_k) \mid i_k \in I, U_k \in \tau_{A_{i_k}}, 1 \leq k \leq n, 0 \leq n\}.$$

Then τ is an open base on S . It is straightforward to see that $f_i : (S, \tau) \rightarrow X_i$ for each $i \in I$, and hence if $h : Y \rightarrow (S, \tau)$, then $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. Let Y be a neighbourhood space and let $h : \underline{Y} \rightarrow S$ be such that $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. If $h(x) \in \bigcap_{k=1}^n f_{i_k}^{-1}(U_k)$, then, since $x \in \bigcap_{k=1}^n (f_{i_k} \circ h)^{-1}(U_k)$, there exists $V \in \tau_Y$ such that $x \in V \subseteq \bigcap_{k=1}^n (f_{i_k} \circ h)^{-1}(U_k) = h^{-1}(\bigcap_{k=1}^n f_{i_k}^{-1}(U_k))$. Hence $h : B \rightarrow (S, \tau)$. \square

Ishihara and Palmgren [15, Theorem 4.3] proved the following proposition in \mathbf{CZF} with the *Relativized Dependent Choice* (RDC).

Proposition 3.2. *Let S be a set, let $\{A_i\}_{i \in I}$ be a family of neighbourhood spaces, and for each $i \in I$ let $f_i : \underline{A_i} \rightarrow S$. Then there exists an open base τ on S such that if h is a mapping from S into the underlying set \underline{B} of a neighbourhood space B , then $h : (S, \tau) \rightarrow B$ if and only if $h \circ f_i : A_i \rightarrow B$ for each $i \in I$.*

Theorem 3.3. *The category \mathbf{Nbh} is complete and cocomplete.*

Proof. Similar to the proof of Theorem 2.5, using Proposition 3.1 and Proposition 3.2. \square

In the following, we shall write, simply, \mathbf{R} for the neighbourhood space \mathbf{R} with the standard open base consisting of open intervals, and $\mathcal{C}(A)$ for $\mathcal{C}(A, \mathbf{R})$.

Let S be a set, and define a relation Φ_S between a function space structure \mathcal{F} on S and $g \in \mathbf{R}^S$ as follows: $\Phi_S(\mathcal{F}, g)$ if and only if for each $x \in S$ and $\epsilon > 0$ there exist $f_1, \dots, f_n \in \mathcal{F}$ ($n \geq 0$) and $\delta > 0$ such that

$$\forall y \in S \left(\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta \implies |g(x) - g(y)| < \epsilon \right).$$

Then a pre-function space X is Φ -closed if

$$\forall g \in \mathbf{R}^X (\Phi_X(\mathcal{F}_X, g) \implies g \in \mathcal{F}_X).$$

Note that if X is a Φ -closed pre-function space, then \mathcal{F}_X contains the constant functions, and the pointwise sum of finitely many functions in \mathcal{F}_X belongs to \mathcal{F}_X . Furthermore, it is straightforward to show that if X is a Φ -closed pre-function space, then X is $\mathcal{C}(\mathbf{R})$ -complete, and uniform limits of functions of \mathcal{F}_X are in \mathcal{F}_X . It was shown in [5, Lemma 1] that if X is a $\mathcal{C}(\mathbf{R})$ -complete pre-function space such that \mathcal{F}_X is closed under finite pointwise sum, then \mathcal{F}_X is closed under finite pointwise product. Therefore Φ -closed pre-function spaces are function spaces in Bishop's sense.

For each set S , the discrete function space on S is Φ -closed, and the pre-function space (S, \mathcal{K}_S) , called the *constant* function space on S , where \mathcal{K}_S is the set of constant functions on S , is Φ -closed. For each Φ -closed pre-function space Y , any mapping $f : \underline{Y} \rightarrow S$ is a function space morphism from Y into the constant function space of S .

In the following, we shall call a Φ -closed pre-function space just a *function space*.

Let \mathbf{Fun} denote the category of pre-function spaces with function spaces as objects and function space morphisms as morphisms. Note that, in \mathbf{Fun} , the initial object is the discrete function space of \emptyset , the terminal objects are the constant function spaces of singletons.

An *adjunction* $\langle F, G, \eta, \varepsilon \rangle$ between categories \mathbf{C} and \mathbf{D} consists of functors $F : \mathbf{C} \rightarrow \mathbf{D}$ and $G : \mathbf{D} \rightarrow \mathbf{C}$, and natural transformations $\eta : 1_{\mathbf{C}} \rightarrow GF$ and $\varepsilon : FG \rightarrow 1_{\mathbf{D}}$ such that $\varepsilon_F \circ F\eta = 1_F$ and $G\varepsilon \circ \eta_G = 1_G$. The functor F is the *left-adjoint*, and the functor G is the *right-adjoint*. The natural transformation η is the *unit*, and the natural transformation ε is the *counit*.

The adjunction $\langle F, G, \eta, \varepsilon \rangle$ is called an *adjoint equivalence* if both the unit η and the counit ε are natural isomorphisms.

We now aim to prove the following result.

Theorem 3.4. *There exists an adjunction between \mathbf{Nbh} and \mathbf{Fun} whose counit is a natural isomorphism.*

Corollary 3.5. *The category \mathbf{Fun} is complete and cocomplete.*

Proof. Since the category \mathbf{Nbh} is complete and cocomplete, by Theorem 3.3, it follows from [16, Exercise 7, VI.3] or the dual of Theorem 4.2 in [14]. \square

To prove Theorem 3.4, we need a series of lemmas.

Lemma 3.6. *For each neighbourhood space A , the pre-function space $(\underline{A}, \mathcal{C}(A))$ is Φ -closed.*

Proof. Let $g \in \mathbf{R}^{\underline{A}}$, and suppose that $\Phi(\mathcal{C}(A), g)$. Then for each $x \in \underline{A}$ and $\epsilon > 0$ there exist $f_1, \dots, f_n \in \mathcal{C}(A)$ and $\delta > 0$ such that for each $y \in \underline{A}$, if $\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta$, then $|g(x) - g(y)| < \epsilon$. Hence there exists $U \in \tau_A$ such that $x \in U$ and if $y \in U$, then $\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta$, and hence $|g(x) - g(y)| < \epsilon$. Thus $g \in \mathcal{C}(A)$. \square

Lemma 3.7. *Let A and B be neighbourhood spaces. If $f : A \rightarrow B$, then $f : (\underline{A}, \mathcal{C}(A)) \rightarrow (\underline{B}, \mathcal{C}(B))$.*

Proof. Straightforward. \square

For a function space structure \mathcal{F} on a set S , let $\tau_{\mathcal{F}}$ be the set of subsets of S of the form

$$U_{f_1, \dots, f_n}(x, \epsilon) = \{y \in S \mid \sum_{k=1}^n |f_k(x) - f_k(y)| < \epsilon\},$$

where $f_1, \dots, f_n \in \mathcal{F}$ ($n \geq 0$), $x \in S$ and $\epsilon > 0$.

Lemma 3.8. *For each pre-function space X , the pair $(\underline{X}, \tau_{\mathcal{F}_X})$ is a neighbourhood space.*

Proof. Straightforward. For (NS1), note that $U(x, \epsilon) = \{y \in \underline{X} \mid 0 < \epsilon\} = \underline{X}$. \square

Lemma 3.9. *Let X and Y be pre-function space. If $f : X \rightarrow Y$, then $f : (\underline{X}, \tau_{\mathcal{F}_X}) \rightarrow (\underline{Y}, \tau_{\mathcal{F}_Y})$.*

Proof. Suppose that $f : X \rightarrow Y$, and let $f(x) \in U_{g_1, \dots, g_n}(y, \epsilon) \in \tau_{\mathcal{F}_Y}$. Then, since $g_i \circ f \in \mathcal{F}_X$ for each $i = 1, \dots, n$, we have $x \in U_{g_1 \circ f, \dots, g_n \circ f}(x, \delta) \in \tau_{\mathcal{F}_X}$ with

$$\delta = \epsilon - \sum_{i=1}^n |g_i(y) - g_i(f(x))|,$$

and if $z \in U_{g_1 \circ f, \dots, g_n \circ f}(x, \delta)$, then $f(z) \in U_{g_1, \dots, g_n}(y, \epsilon)$. Hence $f : (\underline{X}, \tau_{\mathcal{F}_X}) \rightarrow (\underline{Y}, \tau_{\mathcal{F}_Y})$. \square

Lemma 3.10. *If A is a neighbourhood space, then $\text{id}_A : A \rightarrow (\underline{A}, \tau_{\mathcal{C}(A)})$.*

Proof. Let $x \in U_{f_1, \dots, f_n}(y, \epsilon) \in \tau_{\mathcal{C}(A)}$. Then, since $f_1, \dots, f_n \in \mathcal{C}(A)$, there exists $U \in \tau_A$ such that $x \in U$ and if $z \in U$, then

$$\sum_{i=1}^n |f_i(x) - f_i(z)| < \epsilon - \sum_{i=1}^n |f_i(y) - f_i(x)|,$$

and hence $z \in U_{f_1, \dots, f_n}(y, \epsilon)$. Thus $\text{id}_A : A \rightarrow (\underline{A}, \tau_{\mathcal{C}(A)})$. \square

Lemma 3.11. *Let X be a pre-function space. Then $\mathcal{F}_X \subseteq \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X})$. Moreover, if X is Φ -closed, then $\mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X}) \subseteq \mathcal{F}_X$.*

Proof. Note that $g \in \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X})$ if and only if $\Phi_{\underline{X}}(\mathcal{F}_X, g)$. Then, trivially, $\mathcal{F}_X \subseteq \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X})$, and if X is Φ -closed, then $\mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X}) \subseteq \mathcal{F}_X$. \square

Proof of Theorem 3.4. Define a functor F from \mathbf{Nbh} to \mathbf{Fun} by $F(A) = (\underline{A}, \mathcal{C}(A))$ and $F(f) = f$, and define a functor G from \mathbf{Fun} to \mathbf{Nbh} by $G(X) = (\underline{X}, \tau_{\mathcal{F}_X})$ and $G(f) = f$. Then F and G are faithful functors, by Lemma 3.9 and Lemma 3.7.

Furthermore, we see that if we let η_A and ϵ_X denote the identity maps on the sets \underline{A} and \underline{X} , respectively, then $\eta_A : A \rightarrow (\underline{A}, \tau_{\mathcal{C}(A)})$ in \mathbf{Nbh} , by Lemma 3.10, and

$$\epsilon_X : (\underline{X}, \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X})) \rightarrow X \text{ and } \epsilon_X^{-1} : X \rightarrow (\underline{X}, \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X}))$$

in \mathbf{Fun} , by Lemma 3.11. Hence $\eta : 1_{\mathbf{Nbh}} \rightarrow GF$ is a natural transformation and $\epsilon : FG \rightarrow 1_{\mathbf{Fun}}$ is a natural isomorphism satisfying $\epsilon_F \circ F\eta = 1_F$ and $G\epsilon \circ \eta_G = 1_G$. Therefore $\langle F, G, \eta, \epsilon \rangle$ forms an adjunction between \mathbf{Nbh} and \mathbf{Fun} . \square

Let S be a set. Then a family $\{d_i\}_{i \in I}$ of pseudometrics on S is *compatible* with an open base τ on S if

1. for each $x \in S$ and $i \in I$, the mapping $y \mapsto d_i(x, y)$ is in $\mathcal{C}(S, \tau)$,
2. for each $x \in S$ and $U \in \tau$ with $x \in U$ there exist $i_1, \dots, i_n \in I$ ($n \geq 0$) and $\delta > 0$ such that for each $y \in S$, if $\sum_{k=1}^n d_{i_k}(x, y) < \delta$, then $y \in U$.

For a function space structure \mathcal{F} on a set S , let $\{d_f\}_{f \in \mathcal{F}}$ be a family of pseudometrics of S defined by $d_f(x, y) = |f(x) - f(y)|$. Then the family $\{d_f\}_{f \in \mathcal{F}}$ is compatible with $\tau_{\mathcal{F}}$.

A neighbourhood space A has a compatible family of pseudometrics if there exists a family $\{d_i\}_{i \in I}$ of pseudometrics on \underline{A} compatible with τ_A .

Lemma 3.12. *If A is a neighbourhood space having a compatible family of pseudometrics, then $\text{id}_{\underline{A}} : (\underline{A}, \tau_{\mathcal{C}(A)}) \rightarrow A$.*

Proof. Let $\{d_i\}_{i \in I}$ be a family of pseudometrics on \underline{A} compatible with τ_A . Let $x \in \underline{A}$ and let $U \in \tau_A$ with $x \in U$. Then there exist $i_1, \dots, i_n \in I$ and $\delta > 0$ such that if $y \in \underline{A}$ and $\sum_{k=1}^n d_{i_k}(x, y) < \delta$, then $y \in U$. Since the mapping $y \mapsto d_{i_k}(x, y)$ is in $\mathcal{C}(A)$ for each $k = 1, \dots, n$, setting $f_k(y) = d_{i_k}(x, y)$, we have $f_k \in \mathcal{C}(A)$ for each $k = 1, \dots, n$, and if $y \in \underline{A}$ and

$$\sum_{k=1}^n d_{i_k}(x, y) = \sum_{k=1}^n |f_k(x) - f_k(y)| < \delta,$$

then $y \in U$. Hence $\text{id}_{\underline{A}} : (\underline{A}, \tau_{\mathcal{C}(A)}) \rightarrow A$. \square

Let $\mathbf{Nbh}_{\text{pms}}$ denote the category of neighbourhood spaces whose objects are neighbourhood spaces having a compatible family of pseudometrics, and whose morphisms are continuous mappings.

Theorem 3.13. *There exists an adjoint equivalence between $\mathbf{Nbh}_{\text{pms}}$ and \mathbf{Fun} .*

Proof. Let F' be a functor restricting the functor F constructed in the proof of Theorem 3.4 to the category $\mathbf{Nbh}_{\text{pms}}$, and note that the functor G constructed in the proof is a functor from \mathbf{Fun} into $\mathbf{Nbh}_{\text{pms}}$. We see that if we let $\eta_A = \text{id}_{\underline{A}}$ and $\epsilon_X = \text{id}_{\underline{X}}$, then $\eta_A : A \rightarrow (\underline{A}, \tau_{\mathcal{C}(A)})$ and $\eta_A^{-1} : (\underline{A}, \tau_{\mathcal{C}(A)}) \rightarrow A$ in $\mathbf{Nbh}_{\text{pms}}$, by Lemma 3.12, and

$$\epsilon_X : (\underline{X}, \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X})) \rightarrow X \text{ and } \epsilon_X^{-1} : X \rightarrow (\underline{X}, \mathcal{C}(\underline{X}, \tau_{\mathcal{F}_X}))$$

in \mathbf{Fun} . Hence $\eta : 1_{\mathbf{Nbh}} \rightarrow GF'$ and $\epsilon : F'G \rightarrow 1_{\mathbf{Fun}}$ are natural isomorphisms satisfying $\epsilon_{F'} \circ F'\eta = 1_{F'}$ and $G\epsilon \circ \eta_G = 1_G$. Therefore $\langle F', G, \eta, \epsilon \rangle$ forms an adjoint equivalence between $\mathbf{Nbh}_{\text{pms}}$ and \mathbf{Fun} . \square

Corollary 3.14. *The category $\mathbf{Nbh}_{\text{pms}}$ is complete and cocomplete.*

4 Uniform and function spaces

In this paper, we define a notion of a uniform space using a base of uniformity as in [13] which is different from the one in [9] and related papers. A *uniform space* A is pair $(\underline{A}, \mathcal{U}_A)$ consisting of a set \underline{A} and an inhabited set \mathcal{U}_A of subsets of $\underline{A} \times \underline{A}$, called a *uniformity* on \underline{A} , such that

$$\text{Ub1. } \forall U, V \in \mathcal{U}_A \exists W \in \mathcal{U}_A (W \subseteq U \cap V),$$

$$\text{Ub2. } \forall U \in \mathcal{U}_A (\Delta \subseteq U),$$

$$\text{Ub3. } \forall U \in \mathcal{U}_A \exists V \in \mathcal{U}_A (V \subseteq U^{-1}),$$

$$\text{Ub4. } \forall U \in \mathcal{U}_A \exists V \in \mathcal{U}_A (V \circ V \subseteq U).$$

Here $\Delta = \{(x, x) \mid x \in \underline{A}\}$, and $U^{-1} = \{(x, y) \mid (y, x) \in U\}$ and

$$U \circ V = \{(x, z) \mid \exists y ((x, y) \in V \wedge (y, z) \in U)\}$$

for each $U, V \subseteq \underline{A} \times \underline{A}$. A *uniformly continuous mapping* f from a uniform space A into a uniform space B is a mapping $f : \underline{A} \rightarrow \underline{B}$ such that

$$\forall V \in \mathcal{U}_B \exists U \in \mathcal{U}_A \forall x, y \in \underline{A} [(x, y) \in U \implies (f(x), f(y)) \in V].$$

We write $f : A \rightarrow B$ to denote that f is a uniformly continuous mapping from A into B , and $\mathcal{C}_u(A, B)$ for the set of uniformly continuous mappings from A into B .

For any set S , there are the uniform spaces $(S, \{\Delta\})$ and $(S, \{S \times S\})$, called the *discrete* uniform space of S and the *trivial* uniform space of S , respectively. For each uniform space Y , any mapping $f : S \rightarrow \underline{Y}$ is a uniformly continuous mapping from the discrete uniform space of S into Y , and any mapping $f : \underline{Y} \rightarrow S$ is a uniformly continuous mapping from Y into the trivial uniform space of S .

Let **Uni** denote the category of uniform spaces whose objects are uniform spaces and whose morphisms are uniformly continuous mappings. Note that, in **Uni**, the initial object is the discrete uniform space of \emptyset , and the terminal objects are the trivial uniform spaces of singletons.

Proposition 4.1. *Let S be a set, let $\{A_i\}_{i \in I}$ be a family of uniform spaces, and for each $i \in I$ let $f_i : S \rightarrow \underline{A}_i$. Then there exists a uniformity \mathcal{U} on S such that if h is a mapping from the underlying set \underline{B} of a uniform space B into S , then $h : B \rightarrow (S, \mathcal{U})$ if and only if $f_i \circ h : B \rightarrow A_i$ for each $i \in I$.*

Proof. Let

$$\mathcal{U} = \left\{ \bigcap_{k=1}^n (f_{i_k} \times f_{i_k})^{-1}(U_k) \mid i_k \in I, U_k \in \mathcal{U}_{A_{i_k}}, 1 \leq k \leq n, 0 \leq n \right\},$$

where $f_i \times f_i : S \times S \rightarrow \underline{A_i} \times \underline{A_i}$ is a mapping with $(f_i \times f_i)(x, y) = (f_i(x), f_i(y))$. Then \mathcal{U} is a uniformity on S .

It is straightforward to see that $f_i : (S, \mathcal{U}) \rightarrow X_i$ for each $i \in I$, and hence if $h : Y \rightarrow (S, \mathcal{U})$, then $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. Let Y be a uniform space and let $h : \underline{Y} \rightarrow S$ be such that $f_i \circ h : Y \rightarrow X_i$ for each $i \in I$. Then for each $i_1, \dots, i_n \in I$ and $U_1 \in \mathcal{U}_{A_{i_1}}, \dots, U_n \in \mathcal{U}_{A_{i_n}}$, there exists $V \in \mathcal{U}_Y$ such that if $(x, y) \in V$, then $(f_{i_k}(h(x)), f_{i_k}(h(y))) \in U_k$ for each $k = 1, \dots, n$, and hence

$$(h(x), h(y)) \in \bigcap_{k=1}^n (f_{i_k} \times f_{i_k})^{-1}(U_k).$$

Therefore $h : B \rightarrow (S, \mathcal{U})$. □

Theorem 4.2. *The category \mathbf{Uni} is complete.*

Proof. Similar to the proof of Theorem 2.5, using Proposition 4.1. □

In the following, we shall write, simply, \mathbf{R} for the uniform space \mathbf{R} with the standard uniformity, and $\mathcal{C}_u(A)$ for $\mathcal{C}_u(A, \mathbf{R})$.

Let S be a set, and define a relation Ψ_S between a function space structure \mathcal{F} on S and $g \in \mathbf{R}^S$ as follows: $\Psi_S(\mathcal{F}, g)$ if and only if for each $\epsilon > 0$ there exist $f_1, \dots, f_n \in \mathcal{F}$ ($n \geq 0$) and $\delta > 0$ such that

$$\forall x, y \in S \left(\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta \implies |g(x) - g(y)| < \epsilon \right).$$

Then a pre-function space X is Ψ -closed if

$$\forall g \in \mathbf{R}^X (\Psi_X(\mathcal{F}_X, g) \implies g \in \mathcal{F}_X).$$

Note that if a pre-function space is Φ -closed, then it is Ψ -closed. If X is a Ψ -closed pre-function space, then \mathcal{F}_X contains the constant functions, and the pointwise sum of finitely many functions in \mathcal{F}_X belongs to \mathcal{F}_X . Furthermore, it is straightforward to show that if X is a Φ -closed pre-function space, then X is $\mathcal{C}_u(\mathbf{R})$ -complete, and uniform limits of functions of \mathcal{F}_X are in \mathcal{F}_X .

For each set S , the discrete function space and the constant function space are Ψ -closed. For each Ψ -closed pre-function space Y , any mapping $f : \underline{Y} \rightarrow S$ is a function space morphism from Y into the constant function space of S .

In the following, we shall call a Ψ -closed pre-function space a *uniform function space*.

Let \mathbf{Fun}_u denote the category of pre-function spaces with uniform function spaces as objects and function space morphisms as morphisms. Note that, in \mathbf{Fun}_u , the initial object is the discrete function space of \emptyset , the terminal objects are the constant function spaces of singletons.

We now aim to prove the following result.

Theorem 4.3. *There exists an adjunction between \mathbf{Uni} and \mathbf{Fun}_u whose counit is a natural isomorphism.*

Corollary 4.4. *The category \mathbf{Fun}_u is complete.*

To prove Theorem 4.3, we need a series of lemmas.

Lemma 4.5. *For each uniform space A , the pre-function space $(\underline{A}, \mathcal{C}_u(A))$ is Ψ -closed.*

Proof. Let $g \in \mathbf{R}^{\underline{A}}$, and suppose that $\Psi(\mathcal{C}_u(A), g)$. Then for each $\epsilon > 0$ there exist $f_1, \dots, f_n \in \mathcal{C}_u(A)$ and $\delta > 0$ such that for each $x, y \in \underline{A}$, if $\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta$, then $|g(x) - g(y)| < \epsilon$. Hence there exists $U \in \mathcal{U}_A$ such that if $(x, y) \in U$, then $\sum_{i=1}^n |f_i(x) - f_i(y)| < \delta$, and hence $|g(x) - g(y)| < \epsilon$. Thus $g \in \mathcal{C}_u(A)$. \square

Lemma 4.6. *Let A and B be uniform spaces. If $f : A \rightarrow B$, then $f : (\underline{A}, \mathcal{C}_u(A)) \rightarrow (\underline{B}, \mathcal{C}_u(B))$.*

Proof. Straightforward. \square

For a function space structure \mathcal{F} on a set S , let $\mathcal{U}_{\mathcal{F}}$ be the set of subsets of $S \times S$ of the form

$$U_{f_1, \dots, f_n}(\epsilon) = \{(x, y) \in S \times S \mid \sum_{k=1}^n |f_k(x) - f_k(y)| < \epsilon\},$$

where $f_1, \dots, f_n \in \mathcal{F}$ ($n \geq 0$) and $\epsilon > 0$.

Lemma 4.7. For each pre-function space X , the pair $(\underline{X}, \mathcal{U}_{\mathcal{F}_X})$ is a uniform space.

Proof. Straightforward. For (Ub4), note that $U_{f_1, \dots, f_n}(\epsilon/2) \circ U_{f_1, \dots, f_n}(\epsilon/2) \subseteq U_{f_1, \dots, f_n}(\epsilon)$. \square

Lemma 4.8. Let X and Y be pre-function space. If $f : X \rightarrow Y$, then $f : (\underline{X}, \mathcal{U}_{\mathcal{F}_X}) \rightarrow (\underline{Y}, \mathcal{U}_{\mathcal{F}_Y})$.

Proof. Suppose that $f : X \rightarrow Y$ and $U_{g_1, \dots, g_n}(\epsilon) \in \mathcal{U}_{\mathcal{F}_Y}$. Then, since $g_i \circ f \in \mathcal{F}_X$ for each $i = 1, \dots, n$, we have $U_{g_1 \circ f, \dots, g_n \circ f}(\epsilon) \in \mathcal{U}_{\mathcal{F}_X}$, and if $(x, y) \in U_{g_1 \circ f, \dots, g_n \circ f}(\epsilon)$, then $(f(x), f(y)) \in U_{g_1, \dots, g_n}(\epsilon)$. Hence $f : (\underline{X}, \mathcal{U}_{\mathcal{F}_X}) \rightarrow (\underline{Y}, \mathcal{U}_{\mathcal{F}_Y})$. \square

Lemma 4.9. If A is a uniform space, then $\text{id}_A : A \rightarrow (\underline{A}, \mathcal{U}_{\mathcal{C}_u(A)})$.

Proof. Let $U_{f_1, \dots, f_n}(\epsilon) \in \mathcal{U}_{\mathcal{C}_u(A)}$. Then, since $f_1, \dots, f_n \in \mathcal{C}_u(A)$, there exists $U \in \mathcal{U}_A$ such that if $(x, y) \in U$, then $\sum_{i=1}^n |f_i(x) - f_i(y)| < \epsilon$, and hence $(x, y) \in U_{f_1, \dots, f_n}(\epsilon)$. Thus $\text{id}_A : A \rightarrow (\underline{A}, \mathcal{U}_{\mathcal{C}_u(A)})$. \square

Lemma 4.10. Let X be a pre-function space. Then $\mathcal{F}_X \subseteq \mathcal{C}_u(\underline{X}, \mathcal{U}_{\mathcal{F}_X})$. Moreover, if X is Ψ -closed, then $\mathcal{C}_u(\underline{X}, \mathcal{U}_{\mathcal{F}_X}) \subseteq \mathcal{F}_X$.

Proof. Note that $g \in \mathcal{C}_u(\underline{X}, \mathcal{U}_{\mathcal{F}_X})$ if and only if $\Psi_{\underline{X}}(\mathcal{F}_X, g)$. Then, trivially, $\mathcal{F}_X \subseteq \mathcal{C}_u(\underline{X}, \mathcal{U}_{\mathcal{F}_X})$, and if X is Ψ -closed, then $\mathcal{C}_u(\underline{X}, \mathcal{U}_{\mathcal{F}_X}) \subseteq \mathcal{F}_X$. \square

We end with the

Proof of Theorem 4.3. Define a functor F from **Uni** to **Fun_u** by $F(A) = (\underline{A}, \mathcal{C}_u(A))$ and $F(f) = f$, and define a functor G from **Fun** to **Uni** by $G(X) = (\underline{X}, \mathcal{U}_{\mathcal{F}_X})$ and $G(f) = f$. Then F and G are faithful functors, by Lemma 4.8 and Lemma 4.6.

Furthermore, we see that if we let η_A and ϵ_X denote the identity maps on the sets \underline{A} and \underline{X} , respectively, then $\eta_A : A \rightarrow (\underline{A}, \tau_{\mathcal{C}_u(A)})$ in **Uni**, by Lemma 4.9, and

$$\epsilon_X : (\underline{X}, \mathcal{C}_u(\underline{X}, \tau_{\mathcal{F}_X})) \rightarrow X \text{ and } \epsilon_X^{-1} : X \rightarrow (\underline{X}, \mathcal{C}_u(\underline{X}, \tau_{\mathcal{F}_X}))$$

in **Fun_u**, by Lemma 4.10. Hence $\eta : 1_{\mathbf{Uni}} \rightarrow GF$ is a natural transformation and $\epsilon : FG \rightarrow 1_{\mathbf{Fun}_u}$ is a natural isomorphism satisfying $\epsilon_F \circ F\eta = 1_F$ and $G\epsilon \circ \eta_G = 1_G$. Therefore $\langle F, G, \eta, \epsilon \rangle$ forms an adjunction between **Uni** and **Fun_u**. \square

Acknowledgements. The author thanks the Japan Society for the Promotion of Science (Grant-in-Aid for Scientific Research (C) No.19500012) for partly supporting the research.

References

- [1] Peter Aczel, *Aspects of general topology in constructive set theory*, Ann. Pure Appl. Logic **137** (2006), 3–29.
- [2] Peter Aczel and Michael Rathjen, *Notes on constructive set theory*, Report No. 40, Institut Mittag-Leffler, The Royal Swedish Academy of Sciences, 2001.
- [3] Errett Bishop, *Foundations of Constructive Mathematics*, McGraw-Hill, New York, 1967.
- [4] Errett Bishop and Douglas Bridges, *Constructive Analysis*, Springer, Berlin, 1985.
- [5] Douglas Bridges, *Reflections on function spaces*, preprint, University of Canterbury, New Zealand, 2008.
- [6] Douglas Bridges and Fred Richman, *Varieties of Constructive Mathematics*, Cambridge University Press, Cambridge, 1987.
- [7] Douglas Bridges and Luminița Viță, *Apartness spaces as a framework for constructive topology*, Ann. Pure Appl. Logic **119** (2003), 61–8.
- [8] Douglas Bridges and Luminița Viță, *Techniques of Constructive Analysis*, Springer, New York, 2006.
- [9] Douglas Bridges and Luminița Viță, *Apartness and Uniformity – a Constructive Development*, Theory and Applications of Computability, Springer Verlag, Heidelberg, 2011 (to appear).
- [10] Robert Goldblatt, *Topoi*, Second edition, Studies in Logic and the Foundations of Mathematics 98, North-Holland Publishing Co., Amsterdam, 1984.
- [11] Robin J. Grayson, *Concepts of general topology in constructive mathematics and in sheaves I*, Ann. Math. Logic **20** (1981), 1–41.

- [12] Robin J. Grayson, *Concepts of general topology in constructive mathematics and in sheaves II*, Ann. Math. Logic **23** (1982), 55-98.
- [13] Hajime Ishihara, *Two subcategories of apartness spaces*, to appear in Ann. Pure Appl. Logic.
- [14] Hajime Ishihara, Ray Mines, Peter Schuster and Luminița Viță, *Quasi-apartness and neighbourhood spaces*, Ann. Pure Appl. Logic **141** (2006), 296-306.
- [15] Hajime Ishihara and Erik Palmgren, *Quotient topologies in constructive set theory and type theory*, Ann. Pure Appl. Logic **141** (2006), 257-265.
- [16] Saunders Mac Lane, *Categories for the working mathematician*, Second edition, Graduate Texts in Mathematics 5, Springer-Verlag, New York, 1998.
- [17] Saunders Mac Lane and Ieke Moerdijk, *Sheaves in geometry and logic*, Universitext, Springer-Verlag, New York, 1994.
- [18] Colin McLarty, *Elementary Categories, Elementary Toposes*, Oxford University Press, 1996.
- [19] Giovanni Sambin, *Intuitionistic formal spaces – a first communication*, In: D. Skordev ed., *Mathematical logic and its applications*, Plenum, New York, 1987, 187-204.
- [20] Giovanni Sambin, *Some points in formal topology*, Theoret. Comput. Sci. **305** (2003), 347-408.
- [21] Giovanni Sambin, *The Basic Picture*, Oxford Logic Guides, Oxford Univ. Press (to appear).
- [22] Anne S. Troelstra and Dirk van Dalen, *Constructivism in Mathematics*, Vol.I and II, North-Holland, Amsterdam, 1988.

Hajime Ishihara
 School of Information Science
 Japan Advanced Institute of Science and Technology
 Nomi, Ishikawa 923-1292, Japan

Email: ishihara@jaist.ac.jp
Tel: +81-761-51-1206
Fax: +81-761-51-1149