

Title	A note on the sequential version of Λ_1^2 statements
Author(s)	Fujiwara, Makoto; Yokoyama, Keita
Citation	Lecture Notes in Computer Science, 7921: 171-180
Issue Date	2013
Type	Journal Article
Text version	author
URL	http://hdl.handle.net/10119/12142
Rights	This is the author-created version of Springer, Makoto Fujiwara and Keita Yokoyama, Lecture Notes in Computer Science, 7921, 2013, 171-180. The original publication is available at www.springerlink.com , http://dx.doi.org/10.1007/978-3-642-39053-1_20
Description	9th Conference on Computability in Europe, CiE 2013, Milan, Italy, July 1-5, 2013.

A note on the sequential version of Π_2^1 statements

Makoto Fujiwara^{1,2,5} and Keita Yokoyama^{3,4,5}

¹ Mathematical Institute, Tohoku University,
6-3, Aramaki Aoba, Aoba-ku, Sendai, Miyagi, Japan.

² sb0m29@math.tohoku.ac.jp

³ School of Information Science, Japan Advanced Institute of Science and
Technology, Nomi, Ishikawa, Japan,

⁴ y-keita@jaist.ac.jp

Abstract. In connection with uniform computability and intuitionistic provability, the strength of the sequential version of Π_2^1 theorems has been investigated in reverse mathematics. In some examples, we illustrate that it occasionally depends on the way of formalizing the Π_2^1 statement, so the investigation of sequential strength demands careful attention to the formalization. Moreover our results suggest the optimality of Dorais's uniformization theorems.

Keywords: Reverse mathematics, Sequential version, Uniformity, Marriage theorem, Bounded König's lemma, Weak weak König's lemma.

1 Introduction

Definition 1 (Sequential version). *The sequential version of a Π_2^1 statement having the form:*

$$(\spadesuit) \quad \forall X (\varphi(X) \rightarrow \exists Y \psi(X, Y))$$

is the statement

$$\forall \langle X_n \rangle_{n \in \mathbb{N}} (\forall n \varphi(X_n) \rightarrow \exists \langle Y_n \rangle_{n \in \mathbb{N}} \forall n \psi(X_n, Y_n)),$$

where X is possibly a tuple of set (or function) variables. Throughout this paper, we denote the sequential version of a statement T having a form (\spadesuit) as $\text{Seq}(T)$.

Many mathematical statements have the form (\spadesuit) , and their sequential forms have been investigated in order to reveal the lack of uniformity of their proof in classical subsystems of second-order arithmetic (e.g. [2], [3], [6]). For instance,

⁵ The first author is supported by a grant from Shigakukai. The second author is supported by a Japan Society for the Promotion of Science postdoctoral fellowship for young scientists, and by a grant from the John Templeton Foundation.

the intermediate value theorem is provable in RCA_0 , but the sequential version of it is equivalent to WKL (weak König's lemma), and so, not provable in RCA_0 . This is of course caused by the necessity of non-uniformity in the proof in RCA_0 . However, the strength of the sequential version may increase for another reason. In this paper, we concentrate our attention on Π_2^1 statements having the following syntactical form:

$$(\natural')$$

$$\forall X (\exists Z \theta(X, Z) \rightarrow \exists Y \psi(X, Y)),$$

where $\theta(X, Z)$ is arithmetical. Despite the fact that (\natural') is, even in intuitionistic predicate logic, equivalent to the following statement:

$$(\natural)$$

$$\forall X, Z (\theta(X, Z) \rightarrow \exists Y \psi(X, Y)),$$

the sequential version of (\natural') is occasionally stronger than that of (\natural) even if $\theta(X, Z)$ has a very weak complexity such as Π_1^0 . This is caused by the difficulty of obtaining the sequence of Z in (\natural') . Using the finite marriage theorem and the bounded König's lemma, we illustrate this phenomenon. On the other hand, the sequential version of a statement of the form (\natural') is not always stronger than that of (\natural) as we see in the case of the weak weak König's lemma. The important point is that the sequential form of (\natural') captures the difficulty of obtaining a solution Y from X alone while that of (\natural) captures the difficulty of obtaining a solution Y using both X and Z . That is to say, whenever we consider the sequential version of a Π_2^1 statement, we must pay attention to the formalization and what information can be used to obtain a solution.

In addition, it has been recently established in [7] and [1] that the intuitionistic provability of Π_2^1 statements of some syntactical form guarantees its classical sequential provability. Such kind of results are called "uniformization theorems". Our results can be used to show that Dorais's uniformization theorems in [1] are the best possible for the syntactical classes involved.

Throughout this paper, we use the standard notation and terminology in reverse mathematics (cf. [9]). In addition, $\mathbf{x} \subset_{\text{fin}} X$ denotes that \mathbf{x} is a finite subset of X , and \mathbb{Q}^+ denotes the set of positive rational numbers. We recall that $\text{WKL}_0 = \text{RCA}_0 + \text{WKL}$ and $\text{ACA}_0 = \text{RCA}_0 + \text{ACA}$ (arithmetical comprehension).

2 The Finite Marriage Theorem

The so-called marriage theorem for finite graphs states that *a finite binary graph $(B, G; R)$ satisfying the Hall condition:*

$$\forall \mathbf{x} \subset_{\text{fin}} B \exists \mathbf{y} \subset_{\text{fin}} G (|\mathbf{x}| \leq |\mathbf{y}| \wedge \forall g \in \mathbf{y} \exists b \in \mathbf{x} ((b, g) \in R)),$$

has an injection $M \subseteq R$ from B to G . It is well-known that there is a uniform algorithm to construct an injection from a given finite bipartite graph satisfying the Hall condition, which suggests that the sequential version of the finite marriage theorem is provable in RCA_0 . However, it depends on the formalization.

We provide the following two formalizations of the finite marriage theorem.

FMT :

$$\forall B, G, R, k \left(\left(\begin{array}{l} (B, G; R) \text{ is a bipartite graph} \\ \text{which satisfies the Hall condition} \\ \text{and } k \text{ bounds } B \cup G \end{array} \right) \rightarrow \exists M \left(\begin{array}{l} M \subseteq R \\ \text{is injective} \end{array} \right) \right),$$

F'MT :

$$\forall B, G, R \left(\exists k \left(\begin{array}{l} (B, G; R) \text{ is a bipartite graph} \\ \text{which satisfies the Hall condition} \\ \text{and } k \text{ bounds } B \cup G \end{array} \right) \rightarrow \exists M \left(\begin{array}{l} M \subseteq R \\ \text{is injective} \end{array} \right) \right),$$

where “ k bounds $B \cup G$ ” denotes that for all $v \in B \cup G$, $v < k$. Note that the premise of $(\dots \rightarrow \dots)$ in FMT is purely universal. Throughout this paper, we use a little odd notation (e.g. F'MT) to indicate which assumption of uniformity is dropped by sequentializing.

Proposition 2.

1. $\text{RCA}_0 \vdash \text{Seq}(\text{FMT})$.
2. $\text{RCA}_0 \vdash \text{Seq}(\text{F'MT}) \leftrightarrow \text{ACA}$.

Proof. (1) A slight recasting of the proof of the finite marriage theorem in RCA_0 ([4, Theorem 2.1]).

(2) $\text{ACA}_0 \vdash \text{Seq}(\text{F'MT})$ follows from the fact that the infinite marriage theorem is provable in ACA_0 ([5, Theorem 2.2]). For the reverse direction, it suffices to find the range of an injection $f : \mathbb{N} \rightarrow \mathbb{N}$ ([9, Lemma III.1.3]). The basic idea is to construct, simultaneously in RCA_0 , infinite numbers of finite bipartite graphs such that the solution of the i -th graph indicates whether i is in $\text{Rng}(f)$ or not. By Σ_0^0 comprehension, take $\langle B_n \rangle_{n \in \mathbb{N}}$ and $\langle G_n \rangle_{n \in \mathbb{N}}$ as

$$\begin{aligned} b \in B_n &\Leftrightarrow b = 0 \vee f\left(\frac{b-2}{2}\right) = n, \\ g \in G_n &\Leftrightarrow g = 1 \vee f\left(\frac{g-3}{2}\right) = n, \end{aligned}$$

which means that in addition to the underlying sequence $\{0, 1\}_{n \in \mathbb{N}}$ of vertices, the odd numbers are divided into $\{B_n\}_{n \in \mathbb{N}}$ and the even numbers are divided into $\{G_n\}_{n \in \mathbb{N}}$ according to f , and take $\langle R_n \rangle_{n \in \mathbb{N}}$ as

$$\begin{aligned} (b, g) \in R_n &\Leftrightarrow (b, g) = (0, 1) \\ &\vee \left(b = 0 \wedge f\left(\frac{g-3}{2}\right) = n \right) \vee \left(g = 1 \wedge f\left(\frac{b-2}{2}\right) = n \right). \end{aligned}$$

Then it is easy to see that $(B_n, G_n; R_n)$ satisfies the Hall condition for each $n \in \mathbb{N}$. Moreover if n is in the range of f via j , $B_n \cup G_n$ is bounded by $2j + 4$, and otherwise, $B_n \cup G_n$ is bounded by 2. Thus, by $\text{Seq}(\text{F'MT})$, there exists a sequence $\langle M_n \rangle_{n \in \mathbb{N}}$ of injections. Put $S := \{n : M_n(0) \neq 1\}$, then S is the range of f by the above construction. \square

The previous proposition indicates that ACA is not needed to construct an injection from a finite bipartite graph satisfying the Hall condition, and only used to take a sequence of bounds. In fact, the next proposition follows from the previous proposition immediately. (One can even prove it directly.)

Proposition 3. *The following assertion SeqB is equivalent to ACA over RCA₀.*

(SeqB) *For any sequence of sets $\langle X_n \rangle_{n \in \mathbb{N}}$, if X_n is finite for all n , then there exists a function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(n)$ bounds X_n .*

Proof. ACA₀ ⊢ SeqB is straightforward. For the reverse direction, it suffices to show Seq(FMT) from SeqB over RCA₀. Let $\langle (B_n, G_n; R_n) \rangle_{n \in \mathbb{N}}$ be a sequence of finite bipartite graphs satisfying the Hall condition. Using SeqB, we have a function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(n)$ bounds $B_n \cup G_n$ for all $n \in \mathbb{N}$. Then the existence of a sequence of injections follows from Seq(FMT). □

3 The Bounded König's lemma

It is known that the bounded König's lemma, which states that *an infinite tree having a bounding function has an infinite path*, is equivalent to WKL [9, Lemma IV.1.4]. As in the previous section, we provide the two formalizations of it.

$$\text{BKL} : \forall T, g \left(\left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is an infinite tree} \\ \text{and } g : \mathbb{N} \rightarrow \mathbb{N} \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

$$\text{B'KL} : \forall T \left(\exists g \left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is an infinite tree} \\ \text{and } g : \mathbb{N} \rightarrow \mathbb{N} \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

where “ g bounds T ” denotes that for all $\sigma \in T$ and $i < \text{lh}(\sigma)$, $\sigma(i) < g(i)$. In addition, we now treat a weaker version of the bounded König's lemma in which a tree in question is bounded by a constant.

$$\text{B}_c\text{KL} : \forall T, k \left(\left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is an infinite tree} \\ \text{and } k \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

$$\text{B}'_c\text{KL} : \forall T \left(\exists k \left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is an infinite tree} \\ \text{and } k \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

where “ k bounds T ” denotes that for all $\sigma \in T$ and $i < \text{lh}(\sigma)$, $\sigma(i) < k$. Note that the premise of $(\dots \rightarrow \dots)$ in B_cKL is purely universal.

Proposition 4.

1. RCA₀ ⊢ Seq(BKL) ↔ Seq(B_cKL) ↔ WKL.
2. RCA₀ ⊢ Seq(B'KL) ↔ Seq(B'_cKL) ↔ ACA.

Proof. We reason in RCA_0 .

(1) WKL implies $\text{Seq}(\text{WKL})$ ([6, Lemma 5]), and $\text{Seq}(\text{WKL})$ implies $\text{Seq}(\text{BKL})$ by imitating the proof of BKL in WKL_0 ([9, Lemma IV.1.4]). The implication from $\text{Seq}(\text{BKL})$ to $\text{Seq}(\text{B}_c\text{KL})$ is obvious. That from $\text{Seq}(\text{B}_c\text{KL})$ to WKL follows immediately from the fact that binary trees are bounded by 2.

(2) It is straightforward that ACA implies $\text{Seq}(\text{B}'\text{KL})$ by imitating the proof of König's lemma in ACA_0 ([9, Lemma III.7.2]). $\text{Seq}(\text{B}'\text{KL})$ implies $\text{Seq}(\text{B}'_c\text{KL})$. The implication from $\text{Seq}(\text{B}'_c\text{KL})$ to ACA follows from Lemma 11 below. \square

In the reverse mathematics of analysis, the bounded König's lemma corresponds to the Heine/Borel compactness of effectively totally bounded complete separable metric spaces. Thus, to consider the strength of a sequential version of a mathematical statement which is related to Heine/Borel compactness, it is important to check which version of bounded König's lemma is needed. Here, we will consider the maximum principle of continuous functions as an example. The following statement is equivalent to WKL over RCA_0 . (See [9, Section IV].)

(MP) For any f , if f is a continuous function from $[-1, 1]$ to \mathbb{R} , then there exists $a \in [-1, 1]$ such that

$$\max\{f(x) : x \in [-1, 1]\} = f(a).$$

By an easy consideration, we can see that MP is equivalent to the following.

(MP⁺) For any f , if f is a continuous function from $(-1, 1)$ to \mathbb{R} such that $f(0) > 0$ and $\lim_{x \rightarrow \pm 1} f(x) = 0$, then there exists $a \in (-1, 1)$ such that

$$\max\{f(x) : x \in (-1, 1)\} = f(a).$$

For the sequential version of MP, the following is well-known, actually, it is an easy consequence of $\text{RCA}_0 \vdash \text{WKL} \leftrightarrow \text{Seq}(\text{WKL})$ ([6, Lemma 5]).

Proposition 5. *Seq(MP) is equivalent to WKL over RCA_0 .*

However, the sequential version of MP⁺ is strictly stronger than that of MP. (In general, ACA is required to extend a continuous function $f : (-1, 1) \rightarrow \mathbb{R}$ with $\lim_{x \rightarrow \pm 1} f(x) = 0$ into a continuous function from $[-1, 1]$ to \mathbb{R} .)

Proposition 6. *The following are equivalent over RCA_0 .*

1. ACA .
2. *The sequential version of the following statement: for any f , if f is a bounded support continuous function from \mathbb{R} to \mathbb{R} , then there exists $a \in \mathbb{R}$ such that $\max\{f(x) : x \in \mathbb{R}\} = f(a)$. (Here, f is said to have bounded support if there exists $k \in \mathbb{N}$ such that the closure of $\{x \in \mathbb{R} : f(x) \neq 0\}$ is a subset of $[-k, k]$.)*
3. *The sequential version of the following statement: for any f , if f is a continuous function from \mathbb{R} to \mathbb{R} such that $f(0) > 0$ and $\lim_{x \rightarrow \pm\infty} f(x) = 0$, then there exists $a \in \mathbb{R}$ such that $\max\{f(x) : x \in \mathbb{R}\} = f(a)$.*

4. Seq(MP⁺).

Proof. By modifying the proof of $\text{MP} \leftrightarrow \text{WKL}$, we can easily see that 2 is equivalent to the sequential version of the following statement: if $T \subseteq \mathbb{N}^{\mathbb{N}}$ is an infinite tree such that $T \subseteq 2k \times 2^{<\mathbb{N}}$ for some k , then T has an infinite path. Note that this is a weaker version of Seq(B_cKL), and still is equivalent to ACA as in the proof of Lemma 11 below. For a given continuous function f from \mathbb{R} to \mathbb{R} such that $f(0) > 0$ and $\lim_{|x| \rightarrow \infty} f(x) = 0$, define a continuous function g as $g(x) = \max\{0, f(x) - f(0)/2\}$. Then, g has bounded support and $\max\{g(x) : x \in \mathbb{R}\} + f(0)/2 = \max\{f(x) : x \in \mathbb{R}\}$, hence we have $2 \leftrightarrow 3$. By an easy rescaling, we have $3 \leftrightarrow 4$. Thus, they are all equivalent to ACA. \square

4 The Weak Weak König's Lemma

The weak weak König's lemma, which states that *a binary tree with positive measure has an infinite path*, has an intermediate strength between RCA_0 and WKL_0 ([9, Remark X.1.8]). In this case, both of sequential versions are stronger than the instancewise version and actually equivalent to WKL .

$$\text{WWKL} : \forall T, m \left(\left(\begin{array}{l} T \subseteq 2^{<\mathbb{N}} \text{ is a tree and} \\ m \in \mathbb{Q}^+ \text{ satisfies } (W_2) \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

$$\text{W'WKL} : \forall T \left(\exists m \left(\begin{array}{l} T \subseteq 2^{<\mathbb{N}} \text{ is a tree and} \\ m \in \mathbb{Q}^+ \text{ satisfies } (W_2) \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

where (W_2) denotes

$$\lim_{n \rightarrow \infty} \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{2^n} \geq m.$$

Proposition 7.

1. $\text{RCA}_0 \vdash \text{Seq}(\text{WWKL}) \leftrightarrow \text{WKL}$. ([2, Theorem 4.1.(2)])
2. $\text{RCA}_0 \vdash \text{Seq}(\text{W'WKL}) \leftrightarrow \text{WKL}$.

Proof (of 2). It is easy to show within RCA_0 that for binary tree T , if there exists $m \in \mathbb{Q}^+$ such that $\lim_{n \rightarrow \infty} \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{2^n} \geq m$, then T is infinite. Therefore $\text{WKL}_0 \vdash \text{Seq}(\text{W'WKL})$ immediately follows from $\text{WKL}_0 \vdash \text{Seq}(\text{WKL})$ ([6, Lemma 5]). For the reverse direction, $\text{Seq}(\text{W'WKL})$ obviously implies $\text{Seq}(\text{WWKL})$, which is equivalent to WKL over RCA_0 from (1). \square

Remark 8. Note that the previous proposition does not suggest that the sequential strength of a mathematical statement equivalent to WWKL is WKL in general. Here, we will consider Riemann integrability for bounded functions as an example. The following statement is equivalent to WWKL over RCA_0 . (See [8].)

(Int) For any f , if f is a continuous function from $[0, 1]$ to $[0, 1]$, then there exists $r \in \mathbb{R}$ such that

$$\int_0^1 f(x) dx = r.$$

However, $\text{Seq}(\text{Int})$ does not imply WKL. This is because $\text{Seq}(\text{Int})$ follows from the following sequential contrapositive of W'WKL:

$$(\star) \quad \forall T \left(\forall n \left(\begin{array}{l} T_n \subseteq 2^{<\mathbb{N}} \text{ is a tree} \\ \text{which has no path} \end{array} \right) \rightarrow \forall n \lim_{k \rightarrow \infty} \frac{|\{\sigma \in T_n : \text{lh}(\sigma) = k\}|}{2^k} = 0 \right).$$

The contrapositive of W'WKL does not have the form (\spadesuit) from Definition 1 any more and (\star) is trivially equivalent to WWKL. Therefore $\text{Seq}(\text{Int})$ is actually equivalent to WWKL. In fact, for many sequential versions of mathematical statements which are equivalent to WWKL, we do not need $\text{Seq}(\text{WWKL})$ or $\text{Seq}(\text{W'WKL})$ but (\star).

Next, we will investigate the effect of uniformity for positive measure more precisely. For this, we shall consider some more variants, namely, bounded König's lemmas with respect to measure.

$$- \text{WBKL} : \forall T, m, g \left(\left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is a tree,} \\ m \in \mathbb{Q}^+ \text{ satisfies } (W_g), \\ g : \mathbb{N} \rightarrow \mathbb{N} \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

where (W_g) denotes

$$\lim_{n \rightarrow \infty} \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{\prod_{i < n} g(i)} \geq m.$$

$$- \text{WB}_c\text{KL} : \forall T, m, k \left(\left(\begin{array}{l} T \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is a tree,} \\ m \in \mathbb{Q}^+ \text{ satisfies } (W_k), \\ k \text{ bounds } T \end{array} \right) \rightarrow \exists P \left(\begin{array}{l} P \text{ is an infinite} \\ \text{path of } T \end{array} \right) \right),$$

where (W_k) denotes

$$\lim_{n \rightarrow \infty} \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{k^n} \geq m.$$

Proposition 9. WBKL and WB_cKL are equivalent to WWKL over RCA_0 .

Proof. We reason in RCA_0 . WBKL to WB_cKL to WWKL is trivial. We will show WBKL from WWKL. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be a tree bounded by $g : \mathbb{N} \rightarrow \mathbb{N}$ such that for some $q \in \mathbb{Q}^+$,

$$\lim_{n \rightarrow \infty} \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{\prod_{i < n} g(i)} \geq q.$$

For $\sigma \in \mathbb{N}^{<\mathbb{N}}$, define $l_g(\sigma)$ and $r_g(\sigma)$ as follows:

$$l_g(\sigma) = \sum_{k < \text{lh}(\sigma)} \frac{\sigma(k)}{\prod_{i \leq k} g(i)}, \quad r_g(\sigma) = l_g(\sigma) + \frac{1}{\prod_{i < \text{lh}(\sigma)} g(i)}.$$

Similarly, for $\sigma \in 2^{<\mathbb{N}}$, define $l_2(\sigma)$ and $r_2(\sigma)$ as follows:

$$l_2(\sigma) = \sum_{k < \text{lh}(\sigma)} \sigma(k)2^{-k-1}, \quad r_2(\sigma) = l_2(\sigma) + 2^{-\text{lh}(\sigma)}.$$

Note that $\bigcup_{\sigma \in T, \text{lh}(\sigma)=m} [l_g(\sigma), r_g(\sigma)]$ are disjoint intervals in $[0, 1]$ whose lengths sum to the measure of level m of T and these intervals can be approximated arbitrarily well from within by intervals with dyadic rational endpoints. That is, for $m \in \mathbb{N}$, there exists $N \in \mathbb{N}$ such that

$$\begin{aligned} & \left| \frac{\left\{ \sigma \in 2^{<\mathbb{N}} : \text{lh}(\sigma) = N \wedge \exists \tau \in T(\text{lh}(\tau) = m \wedge l_g(\tau) \leq l_2(\sigma) \wedge r_2(\sigma) \leq r_g(\tau)) \right\}}{2^N} \right| \\ & > \frac{|\{\sigma \in T : \text{lh}(\sigma) = m\}|}{\prod_{i < m} g(i)} - \frac{q}{2^{m+2}}. \end{aligned}$$

We define $h(m)$ as the least such N .

Now we define $T^* \subseteq 2^{\mathbb{N}}$ as

$$\begin{aligned} \sigma \in T^* & \Leftrightarrow \\ \forall m < \text{lh}(\sigma) & \left(\begin{array}{l} h(m) \leq \text{lh}(\sigma) \rightarrow \exists \tau \in T(\text{lh}(\tau) = m \wedge \\ l_g(\tau) \leq l_2(\sigma \upharpoonright h(m)) \wedge r_2(\sigma \upharpoonright h(m)) \leq r_g(\tau)) \end{array} \right). \end{aligned}$$

Then, T^* is a tree such that for all $n \in \mathbb{N}$,

$$\frac{|\{\sigma \in T^* : \text{lh}(\sigma) = n\}|}{2^n} > \frac{|\{\sigma \in T : \text{lh}(\sigma) = n\}|}{\prod_{i < n} g(i)} - \sum_{m < n} \frac{q}{2^{m+2}} \geq \frac{q}{2}.$$

Thus, by WWKL, there exists a path P^* through T^* . For any $m \in \mathbb{N}$, there exists a unique $\tau_m \in T$ such that $\text{lh}(\tau_m) = m$ and $l_g(\tau_m) \leq l_2(P \upharpoonright h(m)) \wedge r_2(P \upharpoonright h(m)) \leq r_g(\tau_m)$. Then, $P = \bigcup_{m \in \mathbb{N}} \tau_m$ is a path through T . \square

Next we investigate the sequential strength of the statements in question. The following proposition means that the uniformity for positive-measure does not help to weaken the sequential strength of the bounded König's lemma.

Proposition 10.

1. $\text{Seq}(W'BKL)$, $\text{Seq}(WBKL)$, $\text{Seq}(W'B_cKL)$ and $\text{Seq}(WB_cKL)$ are equivalent to WKL over RCA_0 .
2. $\text{Seq}(W'B'KL)$, $\text{Seq}(WB'KL)$, $\text{Seq}(W'B'_cKL)$ and $\text{Seq}(WB'_cKL)$ are equivalent to ACA over RCA_0 .

Here $WB'KL$, $W'BKL$, $W'B'KL$, WB'_cKL , $W'B_cKL$, and $W'B'_cKL$ are defined in the same manner as before, that is, W' (resp. B' , B'_c) means that the universal quantifier $\forall m$ (resp. $\forall g, \forall k$) is moved into $(\dots \rightarrow \dots)$ as the existential quantifier $\exists m$ (resp. $\exists g, \exists k$).

To show the previous proposition, we first show the following lemma.

Lemma 11. $\text{RCA}_0 \vdash \text{Seq}(\text{WB}'_c\text{KL}) \rightarrow \text{ACA}$, that is, the following statement implies ACA over RCA_0 :

$$\forall \langle T_n \rangle_{n \in \mathbb{N}}, \langle m_n \rangle_{n \in \mathbb{N}} \left(\forall n \exists k \left(\begin{array}{l} T_n \subseteq \mathbb{N}^{<\mathbb{N}} \text{ is a tree,} \\ m_n \in \mathbb{Q}^+ \text{ satisfies } (W_k) \text{ for } T_n, \\ k \text{ bounds } T_n \end{array} \right) \right. \\ \left. \longrightarrow \exists \langle P_n \rangle_{n \in \mathbb{N}} \forall n (P_n \text{ is an infinite path of } T_n) \right).$$

Proof. As in the proof of Proposition 2.(2), we will find the range of an injection $f : \mathbb{N} \rightarrow \mathbb{N}$ ([9, Lemma III.1.3]). By Σ_0^0 comprehension, we take a sequence $\langle T_n \rangle_{n \in \mathbb{N}}$ of trees from the given injection f as

$$\sigma \in T_n \Leftrightarrow \begin{array}{l} \forall i < \text{lh}(\sigma) \left(\sigma(0) = 0 \wedge \sigma(i+1) \leq 1 \wedge f(i) \neq n \right) \vee \\ \exists j < \sigma(0) \left(\forall i < \text{lh}(\sigma) \left(\sigma(i) \leq 2j+1 \wedge f(j) = n \right) \right). \end{array}$$

Then, each $T_n \subseteq \mathbb{N}^{<\mathbb{N}}$ clearly forms a tree. Put $m_n := 1/2$. We need to find a required bound k for each n . For given n , if there exists j such that $f(j) = n$, then put $k := 2j + 2$, and otherwise, put $k := 2$. In either case, we can check that k bounds T_n and $m_n (= 1/2)$ satisfies (W_k) for T_n . Thus, by $\text{Seq}(\text{WB}'_c\text{KL})$, there exists a sequence $\langle P_n \rangle_{n \in \mathbb{N}}$ of paths. Put $S := \{n : P_n(0) \neq 0\}$. It is easy to see that $P_n(0) \neq 0 \Leftrightarrow \exists j (f(j) = n)$, namely, S is the range of f . \square

Proof (of Proposition 10). We reason in RCA_0 .

(1) Each of $\text{Seq}(\text{W}'\text{BKL})$, $\text{Seq}(\text{WBKL})$, $\text{Seq}(\text{W}'\text{B}_c\text{KL})$, $\text{Seq}(\text{WB}_c\text{KL})$ follows from $\text{Seq}(\text{BKL})$, then also from WKL by Proposition 4.(1). On the other hand, each of them implies $\text{Seq}(\text{WWKL})$ which is equivalent to WKL .

(2) Each of $\text{Seq}(\text{W}'\text{B}'\text{KL})$, $\text{Seq}(\text{WB}'\text{KL})$, $\text{Seq}(\text{W}'\text{B}'_c\text{KL})$, $\text{Seq}(\text{WB}'_c\text{KL})$ follows from $\text{Seq}(\text{B}'\text{KL})$, then also from ACA by Proposition 4.(2). On the other hand, each of $\text{Seq}(\text{W}'\text{B}'\text{KL})$, $\text{Seq}(\text{WB}'\text{KL})$, $\text{Seq}(\text{W}'\text{B}'_c\text{KL})$ implies $\text{Seq}(\text{WB}'_c\text{KL})$ and $\text{Seq}(\text{WB}'_c\text{KL})$ implies ACA by Lemma 11. \square

5 The Best Possibility of Dorais's Uniformization Results

The first uniformization theorems are established in [7], which can be applied for Π_2^1 statements of the form (\spadesuit) (from Definition 1) with purely universal φ . Dorais has recently shown other uniformization theorems in second-order setting with function-based language, which can be applied for more Π_2^1 statements.

Proposition 12 (Dorais [1]).

1. For any $T : \forall f (\varphi(f) \rightarrow \exists g \psi(f, g))$ such that $\varphi(f)$ is in N_K and $\psi(f, g)$ is in Γ_K , if $\text{EL} + \text{GC} + \text{CN} \vdash T$, then $\text{RCA} \vdash \text{Seq}(T)$.
2. For any $T : \forall f (\varphi(f) \rightarrow \exists g \psi(f, g))$ such that $\varphi(f)$ is in N_L and $\psi(f, g)$ is in Γ_L , if $\text{EL} + \text{WKL} + \text{GC}_L + \text{CN}_L \vdash T$, then $\text{RCA} + \text{WKL} \vdash \text{Seq}(T)$.

We refer the readers to see [1] for precise definitions of each of the symbols in the previous proposition. In fact, the restriction of ψ to Γ_K and Γ_L is not tight and the interest is only in the possibility of extending N_K and N_L . All purely existential and purely universal formulas are included in N_K . In addition, all formulas of the form $\exists x \leq t \forall z A_{qf}$ are included in N_L . Here we show that N_K and N_L cannot be extended to the class including all formulas of the form $\exists x \forall z A_{qf}$ in Proposition 12.

Suppose that in Proposition 12.(1), N_K can be extended to such a class. Since each purely universal formula in set-based language is translated as a purely universal formula in function-based language by identifying sets with their characteristic functions, the premise of $(\dots \rightarrow \dots)$ in function-based F'MT (intuitionistically equivalent to function-based FMT) has the form $\exists x \forall z A_{qf}$. Then Proposition 2.(2) derives that function-based F'MT is not provable in $\text{EL} + \text{GC} + \text{CN}$. However, it is provable in EL_0 by transforming the proof of the finite marriage theorem in RCA_0 ([4, Theorem 2.1]).

Next we suppose in Proposition 12.(2), N_L can be extended to such a class. As in the previous paragraph, Proposition 4.(2) derives that function-based $\text{B}'_c\text{KL}$ (intuitionistically equivalent to function-based B_cKL) is not provable in $\text{EL} + \text{WKL} + \text{GC}_L + \text{CN}_L$. However, it is provable in $\text{EL}_0 + \text{WKL}$ by transforming the proof of the bounded König's lemma in WKL_0 ([9, Lemma IV.1.4]).

Acknowledgment

We are grateful to Ulrich Kohlenbach and anonymous referees for their helpful comments and suggestions on an earlier version of this paper.

References

1. F. G. Dorais, Classical consequences of continuous choice principles from intuitionistic analysis, *Notre Dame Journal to Formal Logic*, to appear.
2. F. G. Dorais, D. D. Dzharov, J. L. Hirst, J. R. Mileti, P. Shafer, On uniform relationships between combinatorial problems, to appear.
3. F. G. Dorais, J. L. Hirst, P. Shafer, Reverse mathematics, trichotomy, and dichotomy, *Journal of Logic and Analysis* **4(13)**, (2012) 1-14
4. J. R. Hirst, Combinatorics in Subsystems of Second Order Arithmetic, *Ph.D. thesis*, Pennsylvania State University, 1987.
5. J. R. Hirst, Marriage theorems and reverse mathematics, *Contemporary Mathematics* **106** (1990), pp. 181–196.
6. J. R. Hirst, Representations of reals in reverse mathematics, *Bull. Pol. Acad. Sci. Math.* **55** (2007), no.4, pp. 303-316.
7. J. R. Hirst and C. Mummert, Reverse mathematics and uniformity in proofs without excluded middle, *Notre Dame Journal to Formal Logic* **52** (2011), no.2, pp. 149–162.
8. Sam Sanders and Keita Yokoyama, The Dirac delta function in two settings of Reverse Mathematics, *Archive for Mathematical Logic* **51** (2012), pp. 99–121.
9. S. G. Simpson, *Subsystems of Second Order Arithmetic Second Edition*, Association for Symbolic Logic. Cambridge University Press, 2009.