

Title	On the Enumeration of Chequered Tilings in Polygons
Author(s)	Hamanaka, Hiroaki; Horiyama, Takashi; Uehara, Ryuhei
Citation	Bridges 2017 Conference Proceedings: 423-426
Issue Date	2017-07-27
Type	Conference Paper
Text version	publisher
URL	http://hdl.handle.net/10119/15124
Rights	Copyright (C) 2017 Authors. Hiroaki Hamanaka, Takashi Horiyama and Ryuhei Uehara, Bridges 2017 Conference Proceedings, 2017, 423-426. http://archive.bridgesmathart.org/2017/bridges2017-423.html
Description	

On the Enumeration of Chequered Tilings in Polygons

Hiroaki Hamanaka
Hyogo University of Teacher Education
942-1 Shimokume Kato-city,
Hyogo 673-1494, Japan
hammer@hyogo-u.ac.jp

Takashi Horiyama
Saitama University
255 Shimo-Okubo, Sakura,
Saitama 338-8570, Japan
horiyama@al.ics.saitama-u.ac.jp

Ryuhei Uehara
Japan Advanced Institute of Science and Technology
1-1 Asahidai, Nomi,
Ishikawa 923-1292, Japan
uehara@jaist.ac.jp

Abstract

The Tokyo 2020 Olympic and Paralympic games emblems, called ‘harmonized chequered emblems,’ are designed with three kinds of rectangles. We enumerate all such tilings in a dodecagon with a hole.

1 Introduction

Figure 1 illustrates the Tokyo 2020 Olympic and Paralympic games emblems, which are called ‘harmonized chequered emblems.’ They are designed with three kinds of rectangles, where the rectangles are derived from three kinds of rhombuses of unit edge length. As shown in Figure 2, the rectangles are obtained by joining the midpoints of the four sides of each rhombus. The rhombuses in Figure 2 respectively have the angles of (a) 30 and 150 degrees, (b) 60 and 120 degrees, and (c) 90 degree. As illustrated in Figure 3, the emblems can be seen as tilings with the rhombuses of a regular dodecagon with edge length two. We enumerate all tilings of the three rhombuses in a dodecagon, where the dodecagon has a hole in the top or center of the shape.




Figure 1 : *Tokyo 2020 Olympic and Paralympic Games Emblems.*

2 Tiling of Rhombuses

While the dodecagons in Figure 3 have holes, we at first consider the case when a dodecagon has no holes. A rhombus tiling of a $2n$ -gon corresponds to a pseudoline arrangement, where two pseudolines cross in at most




Figure 2: Three rectangles and their surrounding rhombuses of the same edge length.




Figure 3: Tilings of three kinds of rhombuses in a dodecagon.




Figure 4: Pseudoline arrangement corresponding to a rhombus tiling.

one point [1]. For example, suppose that we have a rhombus tiling given by the black lines in Figure 4. By connecting the midpoints of parallel sides of all rhombuses, we can obtain the red pseudolines in the figure, where the labels i_L and i_R ($i = 1, 2, \dots, 6$) indicate the end points of the pseudolines. We can see that (1) two pseudolines i_L and i_R in the arrangement do not cross for any $i = 1, 2, \dots, 6$, (2) two pseudolines i_{ℓ_i} and j_{ℓ_j} cross exactly once for $i \neq j$ and $\ell_i, \ell_j \in \{L, R\}$, (3) there is no point where three (or more) pseudolines cross, and (4) each crossing of two pseudolines corresponds to a rhombus in the tiling.

In our case, we have a hole in the top or center of the dodecagon. We can regard the hole as a special piece, and a rhombus tiling with the hole as a tiling with rhombuses and the special piece in the specified position (i.e., in the top or center of the dodecagon). As shown in Figure 5, the pseudolines may be split into two segments by the special piece. We call such an arrangement as a pseudoline arrangement with a hole. In Figure 5 (a), the dodecagon and the hole share two sides 1_L and 6_L . In this case, we say pseudolines 1_L and 6_L are split into two segments although the segments 1_L and 6_L above the hole have length 0. Also





Figure 5: Pseudoline arrangements with holes.



Theorem 1. *Let S_t and S_{pa} be the sets of all tilings of the rhombuses in a $2n$ -gon with a hole and all arrangements of $2n$ pseudolines with a hole, respectively. Then, there is a bijection from S_t to S_{pa} .*

In our case, we introduce a special bar corresponding to the special piece (i.e., the hole) in the tiling, that reverses the labels of pseudoline segments. For example, Figure 7 illustrates Amidakujies corresponding to the pseudoline arrangements in Figure 5. The bars, except for the blue bars, exchange the labels as in the usual Amidakujies. The blue bold bars connecting 6 lines represent the special bars. The special bar in Figure 7(a) exchange the labels from $1_L, 2_L, 3_L, 4_L, 5_L, 6_L$ to $6_L, 5_L, 4_L, 3_L, 2_L, 1_L$. The special bar in Figure 7(b) exchange the labels from $1_R, 2_R, 3_L, 4_L, 5_R, 6_R$ to $6_R, 5_R, 4_L, 3_L, 2_R, 1_R$. We define special Amidakujies as follows: A special Amidakuji is an Amidakuji with $2n$ lines and a special bar and $3\binom{n}{2}$ (usual) bars, where the $2n$ lines have labels $1_L, 1_R, 2_L, 2_R, \dots, n_L, n_R$, a special bar exchange the labels from $1_{\ell_1}, 2_{\ell_2}, \dots, n_{\ell_n}$ to $n_{\ell_n}, (n-1)_{\ell_{n-1}}, \dots, 1_{\ell_1}$ ($\ell_i \in \{L, R\}$ for $i = 1, 2, \dots, n$), and bars do not exchange the same pair of labels twice, and do not exchange the pairs of i_L and i_R ($i = 1, 2, \dots, n$). Now, we have the following theorem.

425




Figure 7: Amidakujies corresponding to the pseudoline arrangements in Figure 5.

3 Chequered Tiling with a Hole

We have designed a DP (dynamic programming) algorithm for enumerating all special Amidakujies, and then converted them into rhombus tilings with holes. The number of rhombus tilings with a hole in the top and center is 3,357,270 and 539,968, respectively. (We distinguish two tilings even if one is a rotation/reflection of another.) Figure 8 is a partial list of the enumerated chequered tilings with a hole in the center of the dodecagon.




Figure 8: Partial list of enumerated tilings.

References

- [1] O. Bodini, T. Fernique, M. Rao, and E. Rémila, Distance on Rhombus Tilings, *Theoretical Computer Science*, 412 (2011), pp. 4787–4794.
- [2] K. Yamanaka, S. Nakano, Y. Matsui, R. Uehara, and K. Nakada, Efficient enumeration of all ladder lotteries and its application, *Theoretical Computer Science*, 411 (2010), pp. 1714–1722.