

| | |
|--------------|---|
| Title | Recent studies on the proof-theoretic strength of Ramsey's theorem for pairs |
| Author(s) | Yokoyama, Keita |
| Citation | RIMS Kokyuroku (数理解析研究所講究録), 2050: 67-76 |
| Issue Date | 2017-10 |
| Type | Departmental Bulletin Paper |
| Text version | author |
| URL | http://hdl.handle.net/10119/16091 |
| Rights | (c)Author. Keita Yokoyama, RIMS Kokyuroku (数理解析研究所講究録), 2050, 2017, pp.67-76. |
| Description | |

Recent studies on the proof-theoretic strength of Ramsey's theorem for pairs

Keita Yokoyama

Japan Advanced Institute of Science and Technology

email: y-keita@jaist.ac.jp

1 Introduction

Calibrating the strength of Ramsey's theorem is one of the central topics in the study of reverse mathematics. Our target is infinite Ramsey's theorem on \mathbb{N} . Within the second-order arithmetic, we consider Ramsey's theorem for n -tuples and k -colors (RT_k^n) which asserts that every k -coloring of $[\mathbb{N}]^n$ admits an infinite homogeneous subset, and we write RT^n for the statement $\forall k \text{RT}_k^n$.

The strength of Ramsey's theorem was precisely analyzed by means of computability theoretic methods, which led the comparison of Ramsey's theorem with the big five systems in the setting of reverse mathematics. In [15], Jockusch showed that there exists a computable coloring for $[\mathbb{N}]^3$ whose homogeneous set always computes the halting problem. This idea together with a standard proof of Ramsey's theorem is formalized by Simpson [24] within the second-order arithmetic, namely, if $n \geq 3$, Ramsey's theorem for n -tuples is equivalent to ACA_0 . The status of Ramsey's theorem for pairs was open for a long time, until Seetapun [23] proved that RT_2^2 is strictly weaker than ACA_0 over RCA_0 . On the relation between WKL_0 and RT_2^2 , Jockusch [15] showed that WKL_0 does not imply RT_2^2 . The converse direction was very difficult, but finally, Liu [19] showed that $\text{RCA}_0 + \text{RT}_2^2$ does not imply WKL_0 by a clever forcing method. Furthermore, there are numerous studies on Ramsey's theorem for pairs and related combinatorial principles mainly from the view point of computability theory. See Hirschfeldt [12] for a gentle introduction to the reverse mathematics studies for Ramsey's theorem.

In this manuscript, we mainly focus on the proof-theoretic strength of Ramsey's theorem for pairs. There are long series of studies on this topic by various people and various methods. In [14], Hirst showed that RT_2^2 implies the Σ_2^0 -bounding principle ($\text{B}\Sigma_2^0$), and

This work is partially supported by JSPS Grant-in-Aid 16K17640, JSPS Core-to-Core Program (A. Advanced Research Networks), and JAIST Research Grant 2016(Houga).

RT^2 implies $B\Sigma_3^0$ over RCA_0 . On the other hand, Cholak, Jockusch and Slaman [6] showed that $WKL_0 + RT_2^2 + I\Sigma_2^0$ is a Π_1^1 -conservative extension of $I\Sigma_2^0$, and $WKL_0 + RT^2 + I\Sigma_3^0$ is a Π_1^1 -conservative extension of $I\Sigma_3^0$. Thus, the first-order strength of RT_2^2 is in between $B\Sigma_2^0$ and $I\Sigma_2^0$, and the first-order strength of RT^2 is in between $B\Sigma_3^0$ and $I\Sigma_3^0$. After this work, many advanced studies are done to investigate the first-order strength of Ramsey's theorem and related combinatorial principles. One of the most important methods for these studies is adapting computability-theoretic techniques for combinatorial principles in nonstandard models of arithmetic. By this method, Chong, Slaman and Yang [9, 8] analyzed slightly weaker but important combinatorial principles ADS, CAC and SRT_2^2 (see e.g., [12] for these principles), and finally they showed that RT_2^2 does not imply $I\Sigma_2^0$ over RCA_0 in [7]. More recently, Chong, Kreuzer and Yang [unpublished] showed that $WKL_0 + SRT_2^2 + WF(\omega^\omega)$ is Π_3^0 -conservative over $RCA_0 + WF(\omega^\omega)$, where $WF(\omega^\omega)$ asserts the well-foundedness of ω^ω .

Another important approach is calibration of the proof-theoretic strength of variations of the Paris-Harrington principle which is deduced from infinite Ramsey's theorem by using the idea of the ordinal analysis. One of the most important result of this line is a sharp upper bounds for the Paris-Harrington principle by Ketonen and Solovay [16]. More recently, Bovykin/Weiermann [5] showed that indicators defined by Paris's density notion can approach the proof-theoretic strength of various versions of Ramsey's theorem, and by a similar method, the author [28] showed that $RT_k^n + WKL_0^*$ is fairly weak and is a Π_2^0 -conservative extension of RCA_0^* , where RCA_0^* is RCA_0 with only Σ_0^0 -induction and the exponentiation. There are many more studies from this view point, e.g., by Kotlarski, Weiermann, et al. [26, 18, 4].

Here, we will overview the recent results on the exact strength of RT_2^2 and RT^2 , namely, $RT_2^2 + WKL_0$ is a Π_3^0 -conservative extension of RCA_0 , and $RT^2 + WKL_0$ is a Π_1^1 -conservative extension of $RCA_0 + B\Sigma_3^0$. The main tool for the former result is Paris's density notion plus the ordinal analysis, while the latter result is derived by computability-theoretic arguments in nonstandard models.

2 The proof-theoretic strength of RT_2^2

In this section, we see the proof-theoretic strength of Ramsey's theorem for pairs and two colors (RT_2^2) based on [21]. A formula φ is said to be $\tilde{\Pi}_n^0$ if it is of the form $\varphi \equiv \forall X \theta$ where θ is Π_n^0 . The main theorem of this section is the following.

Theorem 2.1 (Patey/Yokoyama). *$WKL_0 + RT_2^2$ is a $\tilde{\Pi}_3^0$ -conservative extension of RCA_0 .*

Recall that $RCA_0 + RT_2^2$ implies $B\Sigma_2^0$ and $RCA_0 + B\Sigma_2^0$ is $\tilde{\Pi}_3^0$ -conservative over $I\Sigma_1^0$. Thus, the theorem says that $I\Sigma_1^0$ is the exact $\tilde{\Pi}_3^0$ -part of $WKL_0 + RT_2^2$. This answers the long-standing open question of determining the Π_2^0 -consequences of RT_2^2 posed e.g., in Seetapun

and Slaman [23, Question 4.4] Cholak, Jockusch and Slaman [6, Question 13.2]. Indeed, one can see that RT_2^2 does not imply the totality of Ackermann function nor the consistency of IS_1^0 . Moreover, one can formalize the proof of this theorem within PRA, which means that $\text{WKL}_0 + \text{RT}_2^2$ is equiconsistent with PRA over PRA.

Now, we overview the idea of the proof. The first step to this theorem is the indicator argument with the density notion introduced by Kirby and Paris [17, 20]

Definition 2.1 (RCA_0). • A finite set $X \subseteq \mathbb{N}$ is said to be *0-dense* if $|X| > \min X$.

• A finite set X is said to be *$m+1$ -dense* if for any $P : [X]^2 \rightarrow 2$, there exists $Y \subseteq X$ which is *m -dense* and *P -homogeneous*.

Note that “ X is *m -dense*” can be expressed by a Σ_0^0 -formula. Let $m\text{PH}_2^2$ be the assertion “for any infinite set $X \subseteq \mathbb{N}$, there exists a finite set $F \subseteq X$ such that X is *m -dense*.” The following theorem is a generalization of the theorem by Bovykin/Weiermann in [5].

Theorem 2.2. $\text{WKL}_0 + \text{RT}_2^2$ is a $\tilde{\Pi}_3^0$ -conservative extension of $\text{RCA}_0 + \{m\text{PH}_2^2 \mid m \in \omega\}$.

Thus, what we need for the main theorem is proving $m\text{PH}_2^2$ within RCA_0 for any $m \in \omega$. For this, we will decompose the density notion by α -largeness notion with ordinals $\alpha < \omega^\omega$. (Here, we use the symbols ω, ω^2, \dots for the internal ordinals.)

Definition 2.2 (RCA_0 , see [11] for the general definition). Let $\alpha < \omega^\omega$.

- If $\alpha = 0$, then any set is said to be α -large.
- If $\alpha = \beta + 1$, then X is said to be α -large if $X \setminus \{\min X\}$ is β -large.
- If $\alpha = \beta + \omega^{n+1}$, then X is said to be α -large if $X \setminus \{\min X\}$ is $(\beta + \omega^n \cdot \min X)$ -large.

Now we will work on finite combinatorics for Ramsey’s theorem based on α -largeness notion. For a given $n \in \omega$, we want to find large enough $m \in \omega$ so that for any ω^m -large set $X \subseteq \mathbb{N}$ and for any coloring $P : [X]^2 \rightarrow 2$, there exists $Y \subseteq X$ which is *P -homogeneous* and ω^n -large. For this, the key notions are “transitivity” and “grouping”.

Definition 2.3 (RCA_0). Let $\alpha, \beta < \omega^\omega$. Let $X \subseteq \mathbb{N}$ and let $P : [X]^2 \rightarrow 2$.

- A set $Y \subseteq X$ is said to be *transitive for P* if for any $x, y, z \in Y$ such that $x < y < z$, $P(x, y) = P(y, z) \rightarrow P(x, y) = P(x, z)$. If X is transitive for P , then P is said to be a transitive coloring on X .
- A sequence of finite sets $\langle F_i \subseteq X \mid i < l \rangle$ is said to be an (α, β) -*grouping for P* if
 - $\forall i < j < l \ F_i < F_j$,

- $\forall i < l$ F_i is α -large,
- for any $H \subseteq_{\text{fin}} \mathbb{N}$, if $H \cap F_i \neq \emptyset$ for any $i < l$, then H is β -large, and,
- $\forall i < j < l \exists c < 2 \forall x \in F_i, \forall y \in F_j P(x, y) = c$.

By transitivity, one can decompose the construction of a homogeneous set into two parts, *i.e.*, first find a large enough transitive subset for a given coloring, and then find a homogeneous set for transitive coloring. The idea of this decomposition is essentially due to Bovykin/Weiermann[5] and Hirschfeldt/Shore[13]. In fact, finding a large homogeneous set for a transitive coloring is much easier than the general case since two homogeneous set can be combined easily by transitivity. On the other hand, constructing a large enough transitive set for a given coloring is harder. For this, we use the idea of grouping. A grouping for P is a family of finite sets such that for any pair of sets from the family, the color between them is fixed. If a family of transitive set forms a grouping, then one can combine them as follows: if a family of finite sets $\langle F_i \subseteq X \mid i < l \rangle$ is a grouping for $P : [X]^2 \rightarrow 2$ such that each of F_i is transitive for P and there is a unified color $c < 2$ such that the color between F_i and F_j is c for any $i < j < l$, then the union $\bigcup_{i < l} F_i$ is transitive for P . By these considerations, we have the following combinatorics.

Lemma 2.3 (RCA_0). *Let $n \in \omega$. Let $X \subseteq_{\text{fin}} \mathbb{N}$ and $\min X > 3$. Then we have the following.*

1. *Ketonen/Solovey[16, Section 6], see also Pelupessy[22]: if X is ω^{n+4} -large, then any coloring $P : [X]^2 \rightarrow n$ has an ω -large homogeneous set.*
2. *If X is ω^{2n+6} -large, then any transitive coloring $P : [X]^2 \rightarrow 2$ has an ω^n -large homogeneous set.*
3. *$\langle F_i \subseteq X \mid i < l \rangle$ is a (ω^n, ω) -grouping for $P : [X]^2 \rightarrow 2$ such that each of F_i is transitive for P and there is a unified color $c < 2$ such that the color between F_i and F_j is c for any $i < j < l$, then the union $\bigcup_{i < l} F_i$ is transitive for P which is ω^{n+1} -large.*

The last piece of the proof is the bound for grouping.

Theorem 2.4. *For any $n, k \in \omega$, there exists $m \in \omega$ such that RCA_0 proves the following:*

if $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^m -large and $\min X > 3$, then, for any coloring $P : [X]^2 \rightarrow 2$, there exists an (ω^n, ω^k) -grouping for P .

In [21], this theorem is proved by considering the infinite version of grouping. Indeed, the existence of a large enough finite set which admits finite grouping for any coloring is an easy consequence of the infinite grouping principle, and the infinite grouping principle is

$\tilde{\Pi}_3^0$ -conservative over RCA_0 , which is shown by a variant of Mathias forcing introduced by Cholak/Jockusch/Slaman[6] and the resplendency argument by Barwise/Schlipf[2]. Recently, the theorem is reproved with a more direct method by Kołodziejczyk, Wong and the author.

Proof of Theorem 2.1. By Theorem 2.2, we only need to show that RCA_0 proves that any infinite set contains m -dense finite set for each $m \in \omega$.

In what follows, we only consider finite sets with their minimum greater than 3. We first show by induction that for any $n \in \omega$, there exists $m \in \omega$ such that RCA_0 proves that if a finite set $X \subseteq \mathbb{N}$ is ω^m -large, then any coloring on X has an ω^n -large transitive set. For the case $n = 1$, $m = 6$ is enough by Lemma 2.3.1. Assume now $n > 1$ and any coloring on an ω^{m_0} -large finite set has an ω^{n-1} -large transitive set. By Theorem 2.4, take $m \in \omega$ so that RCA_0 proves any coloring on an ω^m -large finite set has an (ω^{m_0}, ω^6) -grouping. Let $X \subseteq \mathbb{N}$ be ω^m -large, P be a coloring on X , and $\langle F_i \subseteq X \mid i < l \rangle$ be an (ω^{m_0}, ω^6) -grouping for P . Since each F_i is ω^{m_0} -large, there exists $H_i \subseteq F_i$ such that H_i is an ω^{m-1} -large transitive set for P . On the other hand, $\{\max F_i \mid i < l\}$ is ω^6 -large, thus, there exists $\tilde{H} \subseteq \{\max F_i \mid i < l\}$ such that \tilde{H} is ω -large and P is constant on $[\tilde{H}]^2$ by Lemma 2.3.1. Then, by Lemma 2.3.3, $H = \bigcup \{H_i \mid i < l, \max F_i \in \tilde{H}\}$ is an ω^n -large transitive set for P .

Now we see that for any $n \in \omega$, there exists $m \in \omega$ such that RCA_0 proves that if a finite set $X \subseteq \mathbb{N}$ is ω^m -large, then any coloring on X has an ω^n -large homogeneous set. This is an easy consequence of the above claim and Lemma 2.3.2. Thus, by induction, for any $n \in \omega$, there exists $m \in \omega$ such that RCA_0 proves that any ω^m -large finite set is n -dense. Finally, one can easily show that any infinite set contains ω^m -large finite subset for each $m \in \omega$ within RCA_0 . \square

3 The proof-theoretic strength of RT^2

In this section, we see the proof-theoretic strength of Ramsey's theorem for pairs and finitely many colors. Here, we write RT^2 for $\forall k \text{RT}_k^2$. The full version of the proof for the following theorem will be available in [25].

Theorem 3.1 (Slaman/Yokoyama). $\text{WKL}_0 + \text{RT}^2$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_3^0$.

Since $\text{RCA}_0 + \text{RT}^2$ implies $\text{B}\Sigma_3^0$, $\text{B}\Sigma_3^0$ is the exact Π_1^1 -part of $\text{WKL}_0 + \text{RT}^2$. Note that $\text{B}\Sigma_3^0$ is Π_4^0 -conservative over $\text{I}\Sigma_2^0$. Thus, the proof-theoretic strength of $\text{WKL}_0 + \text{RT}^2$ is the same as $\text{I}\Sigma_2^0$. In addition, the proof of this theorem is again formalizable within PRA, and $\text{WKL}_0 + \text{RT}^2$ is equiconsistent with $\text{I}\Sigma_2^0$ over PRA.

The first step of the proof is the standard decomposition of RT^2 by the cohesiveness principle.

Theorem 3.2 (Cholak/Jockusch/Slaman[6]). *Over RCA_0 , RT^2 is equivalent to COH plus D^2 , where,*

- COH : for any sequence of sets $\langle R_n \subseteq \mathbb{N} \mid n \in \mathbb{N} \rangle$, there exists an infinite set $X \subseteq \mathbb{N}$ such that $\forall n (X \subseteq^* R_n \vee X \subseteq^* R_n^c)$.
- D^2 : for every Δ_2^0 -partition $\bigsqcup_{i < k} \mathcal{A}_i = \mathbb{N}$, there exists an infinite set $X \subseteq \mathbb{N}$ such that $X \subseteq \mathcal{A}_i$ for some $i < k$.

For Π_2^1 -theories, two Π_1^1 -conservative extensions can be amalgamated, *i.e.*, for given Π_2^1 -theories T_0, T_1, T_2 , if T_1 and T_2 are Π_1^1 -conservative extensions of T_0 , then $T_1 + T_2$ is also Π_1^1 -conservative over T_0 (see [27]). Thus, we only need to check the conservation for WKL_0 , COH and D^2 independently. A general conservation theorem for WKL_0 and COH over $\text{RCA}_0 + \text{B}\Sigma_n^0$ are calibrated by Hájek[10] and Belanger[3], respectively.

Theorem 3.3 (Hájek, Belanger). *$\text{WKL}_0 + \text{COH} + \text{B}\Sigma_3^0$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_3^0$.*

To obtain a conservation result for D^2 , we will use the basis theorem for RT^2 from the computability theoretic view point.

Theorem 3.4 (Cholak/Jockusch/Slaman[6]). *For every Δ_2^0 -partition $\bigsqcup_{i < k} \mathcal{A}_i = \omega$, there exists an infinite low₂ set $X \subseteq \omega$ such that $X \subseteq \mathcal{A}_i$ for some $i < k$.*

Here, a set $X \subseteq \omega$ is said to be low₂ if $X'' = \mathbf{0}''$. If X is low₂, then Σ_3^0 predicate relative to X is just Σ_3^0 , thus X preserves $\text{B}\Sigma_3^0$. Therefore, if the above theorem is formalizable within $\text{RCA}_0 + \text{B}\Sigma_3^0$, one can obtain a definable solution for each instance of D^2 which preserves $\text{B}\Sigma_3^0$. This is actually possible, but not directly. Here, we will work within a nonstandard model $(M, S) \models \text{B}\Sigma_3^0$, and consider a Δ_2^0 -partition $\bigsqcup_{i < k} \mathcal{A}_i = M$ for some $k \in M$.

The first obstruction is that to construct a low₂ set, we essentially use $\mathbf{0}''$ -primitive recursion, which requires $\text{I}\Sigma_3^0$, but we only have $\text{B}\Sigma_3^0$. To prove Theorem 3.4, one constructs an approximation of a solution $G_0 \subseteq G_1 \subseteq \dots$, and at each stage, decides one Σ_2^0 -formula $\psi_e(G)$ by using the idea of Mathias forcing. However, because of the lack of $\text{I}\Sigma_3^0$ in M , the construction stages may not cover the whole M , *i.e.*, $\{j \mid G_j \text{ exists}\}$ would form a proper Σ_3^0 -cut of M . To overcome this situation, we can use Shore blocking argument, namely, we will decide finitely many Σ_2^0 -formulas up to the use of the previous stage. Then, one can decide all Σ_2^0 -formulas before the construction ends.

Another obstruction is an essential use of Σ_3^0 -least number principle. In the original construction, one would first try constructing the solution on color 0, and if it fails, then

try color 1 with using the information from the previous failure, and repeat this process. However, without IS_3^0 , one cannot repeat this for arbitrary many colors until the construction works since the number of color may be nonstandard. Thus, we have to construct possible solutions for all colors simultaneously. Then, BS_3^0 is just enough to guarantee that the construction works for at least one color.

Formalizing these ideas, we have the following.

Theorem 3.5. *For any $(M; X) \models \text{BS}_3^0$ and for every Δ_2^X -partition $\bigsqcup_{i < k} \mathcal{A}_i = M$, there exists an unbounded set $G \subseteq M$ which is Δ_3^X -definable in $(M; X)$ such that $G \subseteq \mathcal{A}_i$ for some $i < k$, and $(M; X, G) \models \text{BS}_3^0$.*

Now, starting from a model $(M; X) \models \text{BS}_3^0$, one can obtain $S \subseteq \mathcal{P}(M)$ with $X \in S$ such that $(M, S) \models \text{RCA}_0 + \text{D}^2 + \text{BS}_3^0$ by using the above theorem repeatedly. Thus, we have the following.

Corollary 3.6. *$\text{RCA}_0 + \text{D}^2$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{BS}_3^0$.*

Therefore, by the amalgamation of the conservation theorem mentioned above, we have Theorem 3.1.

4 Further studies

About the proof-theoretic/first-order strength of Ramsey's theorem for pairs, there are several more important questions to be considered.

4.1 The first-order part of RT_2^2

By Theorem 3.1, we already know that the first-order part of $\text{WKL}_0 + \text{RT}^2$ is BS_3^0 , but we still don't know what the first-order part of $\text{WKL}_0 + \text{RT}_2^2$ is.

Question 4.1. Is $\text{WKL}_0 + \text{RT}_2^2$ a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$?

Since $\text{WKL}_0 + \text{RT}_2^2$ implies BS_2^0 , BS_2^0 is the weakest possible system which may be the first-order part of $\text{WKL}_0 + \text{RT}_2^2$. To prove Π_1^1 -conservation, we usually consider the following version of ω -extension property.

Question 4.2. For given $(M, S) \models \text{RCA}_0 + \text{BS}_2^0$ and $X \in S$, is there $\bar{S} \subseteq \mathcal{P}(M)$ such that $X \in \bar{S}$ and $(M, \bar{S}) \models \text{WKL}_0 + \text{RT}_2^2$?

One may assume that (M, S) is a countable recursively saturated model. Unfortunately, our proof of Theorem 2.1 does not provide any information about the possibility of the existence of such extension. On the other hand, one can generalize Theorem 2.2 and obtain a characterization for the $\tilde{\Pi}_4^0$ -part of $\text{WKL}_0 + \text{RT}_2^2$.

Definition 4.3 (RCA_0). Let $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\lim_{x \rightarrow \infty} f(x) = \infty$. Then, we define the notion of f - m -density as follows.

- A finite set X is said to be f -0-dense if $|X| > \min X$.
- A finite set X is said to be f - $m + 1$ -dense if for any coloring $P : [X]^2 \rightarrow 2$, there exists a P -homogeneous set $Y \subseteq X$ such that Y is f - m -dense and for any $x \in [\min Y, \max Y]$, $f(x) > \min X$.

As same as the usual density notion, “ X is f - m -dense” can be expressed by a Σ_0^0 formula. Let $m\text{PH}_2^{2+}$ be the assertion “for any $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\lim_{x \rightarrow \infty} f(x) = \infty$, and for any infinite set $X \subseteq \mathbb{N}$, there exists a finite set $F \subseteq X$ such that F is f - m -dense.” Then we have a modification of Theorem 2.2 as follows.

Theorem 4.1. $\text{WKL}_0 + \text{RT}_2^2$ is a $\tilde{\Pi}_4^0$ -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \{m\text{PH}_2^{2+} \mid m \in \omega\}$.

Question 4.4. Is $m\text{PH}_2^{2+}$ provable within $\text{RCA}_0 + \text{B}\Sigma_2^0$ for any $m \in \omega$?

If the answer is positive, then we know that $\text{WKL}_0 + \text{RT}_2^2$ is $\tilde{\Pi}_4^0$ -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.

4.2 Feasibility of the conservation results

Our conservation results are proved by model theoretic arguments. Unfortunately, that doesn’t mean any feasibility of the conservation. For example, if we have a proof for a $\tilde{\Pi}_3^0$ -sentence ψ from $\text{WKL}_0 + \text{RT}_2^2$, then can we find a proof for ψ from RCA_0 in a feasible way? Formally, we can ask the following.

Question 4.5. Is there a polynomial proof transformation for the $\tilde{\Pi}_3^0$ -conservation between RCA_0 and $\text{WKL}_0 + \text{RT}_2^2$?

Question 4.6. Is there a polynomial proof transformation for the Π_1^1 -conservation between $\text{RCA}_0 + \text{B}\Sigma_3^0$ and $\text{WKL}_0 + \text{RT}^2$?

For the latter case, it is actually not so difficult to find a polynomial proof transformation. By the proof of Theorem 3.5, there is a canonical way to construct a Δ_3^0 -definable solution for RT^2 which preserves $\text{B}\Sigma_3^0$ within $\text{RCA}_0 + \text{B}\Sigma_3^0$. Thus, one can always use the solution for RT^2 within $\text{RCA}_0 + \text{B}\Sigma_3^0$ as if RT^2 is available, and WKL is also available within $\text{RCA}_0 + \text{B}\Sigma_3^0$ in a similar way (see [10]). This idea provides a direct interpretation of RT^2 within $\text{RCA}_0 + \text{B}\Sigma_3^0$.

For Question 4.5, the situation is more complicated. Our proof of Theorem 2.1 depends on the indicator argument, which essentially uses a nonstandard model and its initial segment which is not definable in the ground model, but in general, the use of nonstandard

models may bring some conservation result with a super-exponential speed-up. Recently, Kołodziejczyk, Wong and the author studied this question and obtained a reformulation of the indicator argument by means of forcing. Generally speaking, if a model construction for a conservation theorem is provided by forcing, then one would often obtain a polynomial proof transformation as in Avigad[1]. In our case, a canonical polynomial proof transformation for the conservation between RCA_0 and $\text{WKL}_0 + \text{RT}_2^2$ is available by a combination of forcing for the indicator argument plus quantitative proof for Theorem 2.4. Consequently, feasible versions of the conservation results are available in both cases.

References

- [1] Jeremy Avigad. Formalizing forcing arguments in subsystems of second-order arithmetic. *Annals of Pure and Applied Logic*, 82:165–191, 1996.
- [2] John Barwise and John Schlipf. An introduction to recursively saturated and resplendent models. *Journal of Symbolic Logic*, 41(2):531–536, 1976.
- [3] David A. Belanger. Conservation theorems for the cohesiveness principle, 2015. To appear. Available at <http://www.math.nus.edu.sg/~imsdrb/papers/coh-2015-09-30.pdf>.
- [4] Teresa Bigorajska and Henryk Kotlarski. Partitioning α -large sets: some lower bounds. *Trans. Amer. Math. Soc.*, 358(11):4981–5001, 2006.
- [5] Andrey Bovykin and Andreas Weiermann. The strength of infinitary Ramseyan principles can be accessed by their densities. accepted for publication in *Ann. Pure Appl. Logic*, <http://logic.pdmi.ras.ru/~andrey/research.html>, 2005.
- [6] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey’s theorem for pairs. *The Journal of Symbolic Logic*, 66(1):1–15, 2001.
- [7] C. T. Chong, Theodore A. Slaman, and Yue Yang. The inductive strength of Ramsey’s theorem for pairs, 2014. Preprint.
- [8] C. T. Chong, Theodore A. Slaman, and Yue Yang. The metamathematics of Stable Ramsey’s Theorem for Pairs. *J. Amer. Math. Soc.*, 27(3):863–892, 2014.
- [9] Chi-Tat Chong, Theodore A. Slaman, and Yue Yang. Π_1^1 -conservation of combinatorial principles weaker than Ramsey’s Theorem for pairs. *Advances in Matheamtics*, 230:1060–1077, 2012.
- [10] Petr Hájek. Interpretability and fragments of arithmetic. In P. Clote and J. Krajíček, editors, *Arithmetic, Proof Theory and Computational Complexity*, pages 185–196. Oxford, Clarendon Press, 1993.
- [11] Petr Hájek and Pavel Pudlák. *Metamathematics of First-Order Arithmetic*. Springer-Verlag, Berlin, 1993. XIV+460 pages.
- [12] Denis R Hirschfeldt. Slicing the truth. *Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore*, 28, 2014.

- [13] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey's theorem for pairs. *Journal of Symbolic Logic*, 72:171–206, 2007.
- [14] Jeffrey Lynn Hirst. *Combinatorics in Subsystems of Second Order Arithmetic*. PhD thesis, The Pennsylvania State University, August 1987.
- [15] Carl G Jockusch. Ramsey's theorem and recursion theory. *Journal of Symbolic Logic*, 37(2):268–280, 1972.
- [16] Jussi Ketonen and Robert Solovay. Rapidly growing Ramsey functions. *Ann. of Math. (2)*, 113(2):267–314, 1981.
- [17] L. A. S. Kirby and J. B. Paris. Initial segments of models of Peano's axioms. In *Set theory and hierarchy theory V (Proc. Third Conf., Bierutowice, 1976)*, volume 619 of *Lecture Notes in Mathematics*, pages 211–226, 1977.
- [18] Henryk Kotlarski, Bożena Piekart, and Andreas Weiermann. More on lower bounds for partitioning α -large sets. *Annals of Pure and Applied Logic*, 147:113–126, 2007.
- [19] Jiayi Liu. RT_2^2 does not imply WKL_0 . *Journal of Symbolic Logic*, 77(2):609–620, 2012.
- [20] J. B. Paris. Some independence results for Peano Arithmetic. *Journal of Symbolic Logic*, 43(4):725–731, 1978.
- [21] Ludovic Patey and Keita Yokoyama. The proof-theoretic strength of Ramsey's theorem for pairs and two colors. Submitted, 32 pages, available at <http://arxiv.org/abs/1601.00050>.
- [22] Florian Pelupessy. On α -largeness and the Paris-Harrington principle in RCA and RCA_0^* . *ArXiv e-prints*, November 2016.
- [23] David Seetapun and Theodore A. Slaman. On the strength of Ramsey's theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995.
- [24] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Mathematical Logic. Springer-Verlag, 1999. XIV + 445 pages; Second Edition, Perspectives in Logic, Association for Symbolic Logic, Cambridge University Press, 2009, XVI+ 444 pages.
- [25] Theodore A. Slaman and Keita Yokoyama. The strength of ramsey's theorem for pairs and arbitrary many colors. in preparation.
- [26] Andreas Weiermann. A classification of rapidly growing ramsey functions. *Proc. Amer. Math. Soc.*, 132(2):553–561, 2004.
- [27] Keita Yokoyama. On Π_1^1 conservativity of Π_2^1 theories in second order arithmetic. In C. T. Chong et al., editor, *Proceedings of the 10th Asian Logic Conference*, pages 375–386. World Scientific, 2009.
- [28] Keita Yokoyama. On the strength of Ramsey's theorem without Σ_1 -induction. *Math. Log. Q.*, 59(1-2):108–111, 2013.