JAIST Repository

https://dspace.jaist.ac.jp/

Title	Photochemical control of membrane raft organization
Author(s)	Hamada, Tsutomu; Sugimoto, Ryoko; Nagasaki, Takeshi; Takagi, Masahiro
Citation	Soft Matter, 7: 220-224
Issue Date	2011
Туре	Journal Article
Text version	author
URL	http://hdl.handle.net/10119/9875
Rights	Copyright (C) 2011 Royal Society of Chemistry. Tsutomu Hamada, Ryoko Sugimoto, Takeshi Nagasaki, and Masahiro Takagi, Soft Matter, 7, 2011, 220- 224. http://dx.doi.org/10.1039/COSM00797H - Reproduced by permission of The Royal Society of Chemistry
Description	



Japan Advanced Institute of Science and Technology

Photochemical control of membrane raft organization

Tsutomu Hamada,*^a Ryoko Sugimoto,^a Takeshi Nagasaki^b and Masahiro Takagi*^a

Received (in XXX, XXX) Xth XXXXXXXX 200X, Accepted Xth XXXXXXXX 200X First published on the web Xth XXXXXXXX 200X

5 DOI: 10.1039/b00000x

Controllable membrane phase separation through the action of a synthetic photoresponsive amphiphile is reported. We studied multi-component giant vesicles formed from a ternary lipid mixture of saturated and unsaturated phospholipid and cholesterol together with the photoresponsive amphiphile. A change in the conformation of the photoresponsive amphiphile can ¹⁰ switch membrane lateral segregation in a reversible manner. *Cis*-isomerization induces lateral

phase separation in one-phase membranes or produces additional lateral domains in two-phase membranes. Membranes that are close to miscibility boundary show high photo-responsiveness. This is the first report on the reversible control of membrane lateral segregation triggered by a conformational change in a membrane-constituting molecule. These findings may lead to new

¹⁵ methods for controlling membrane self-organization such as raft engineering.

Introduction

The regulation of lateral organization in plasma membranes is essential for physiological functions.¹ Within fluid membranes, microdomains called lipid rafts, which are composed largely 20 of cholesterol and saturated lipids, are laterally segregated

- with high lipid order and slow dynamics.² During molecularassociating events at a membrane interface, such as signaling and trafficking, stabilized microdomains are temporally produced to effectively concentrate specific molecules.³ It is
- ²⁵ hypothesized that raft-stabilizing proteins such as glycosylphosphatidylinositol-anchored proteins (GPI-APs) promote the formation of microdomains.⁴ However, the mechanism of the molecular triggering of microdomain formation is still unclear.
- ³⁰ Membrane lateral segregations are considered to be a form of liquid-order (L_o) - liquid-disorder (L_d) phase separation that develops due to the interaction between lipid molecules; L_o and L_d phases correspond to rafts and the surrounding fluid membrane, respectively.¹ To reveal the physicochemical
- ³⁵ properties of membrane microdomains, studies on model membrane systems using artificial lipid vesicles have been performed.⁵ Recently, with regard to the molecular mechanism for stabilizing microdomains, it has been suggested that subtle changes in membrane components at the
- ⁴⁰ molecular level, induced by protein binding,⁶ actin polymerization,⁷ photoresponsive cholesterol,⁸ and lipid peroxidation,⁹ can affect membrane lateral segregation. Hammond et al., reported that clustering GM₁ with cholera toxin B (CTB) increased miscibility transition temperature to
- ⁴⁵ cause a homogeneous membrane to phase separate into domains.⁶ Liu and Fletcher also reported that actin cytoskeleton through a phosphatidylinositol 4,5 bisphosphate (PIP2) - neural Wiskott-Aldrich syndrome protein (N-WASP) link elevated miscibility temperature to stabilize phase-
- ⁵⁰ separated domains.⁷ Yasuhara et al., observed changes in domain structures under *cis*-isomerization of an azobenzenemodified cholesterol.⁸ Ayuyan and Cohen reported that

photooxidation promoted membrane phase separation.9

- Previously, we reported that a synthetic photoresponsive ⁵⁵ amphiphile (KAON12), the molecular conformation of which can be switched by light irradiation (Scheme 1), can alter the mechanical properties of lipid bilayers to induce shape transitions of homogeneous vesicles, such as budding and hole opening.¹⁰ In the present study, we used KAON12 to achieve ⁶⁰ the controllable formation of lateral domains within heterogeneous membranes. We studied multi-component giant vesicles formed from a ternary lipid mixture of saturated and unsaturated phospholipid and cholesterol together with this photoresponsive amphiphile. To the best of our knowledge,
- 65 this is the first report on the reversible control of membrane lateral segregation triggered by a conformational change in a membrane-constituting molecule.



Scheme 1. Chemical formula of KAON12.

Results

Lateral membrane phase separation

75 Vesicles were prepared from a lipid mixture of the

photoresponsive amphiphile KAON12, dioleoyl phosphatidylcholine (DOPC), dipalmitoyl phosphatidylcholine (DPPC) and cholesterol (Chol). The ternary system of DOPC/DPPC/Chol shows phase separation 5 between a L_o phase composed largely of DPPC and Chol, and a L_d phase which is rich in DOPC. 5 The membranes contain

- 1% of the fluorescent phospholipid rhodamine red-X dihexadecanoyl phosphoethanolamine (rho-PE), which preferentially partitions into the L_d phase.⁵ Figure 1a and b
- ¹⁰ show typical microscopic images of one-phase and two-phase vesicles, respectively, with a composition of 60 mol % lipids (DOPC:DPPC:Chol = 24:12:24) and 40 mol % KAON12. The isomer of KAON12 is the *trans* form. Figure 1c shows the probability of the one-phase and two-phase vesicles as a
- ¹⁵ function of the DOPC:DPPC ratio under a constant fraction of Chol:KAON12=24:40. Membranes with a higher DPPC ratio tend to exhibit phase separation into domains. The coexistence of one-phase and two-phase vesicles at DOPC:DPPC=21:15 is probably due to a wide variation in the mixed fraction of
- $_{\rm 20}$ lipids during the vesicle preparation. We consider that KAON12 is rich in the L_d region similar to DOPC, since it has been previously reported that binary DOPC/KAON12 membranes do not show phase separation. $^{\rm 10}$



²⁵ Fig. 1. Typical microscopic images of one-phase (a) and two-phase (b) vesicles of DOPC/DPPC/Chol/KAON12. (c) Probability of one-phase and two-phase vesicles as a function of the DOPC:DPPC ratio, with 24% Chol and 40% *trans*-KAON12. The number of vesicles observed for each DOPC:DPPC ratio are N = 36 (30:6), 50 (27:9), 90 (24:12), 81 (21:15), 30 85 (18:18), 43 (15:21), and 39 (12:24). White and gray boxes indicate

Photocontrolled membrane phase organization

one-phase and two-phase vesicles, respectively.

First, we investigated the effect of photoirradition on onephase membranes with *trans*-KAON12, and found a photoinduced groups is a particular photo-

- ³⁵ induced reversible change in membrane phase separation. Figure 2a shows a typical micrograph of the reversible molecular switching of membrane phase organization (also see a movie and pictures of other giant vesicles in the Supplementary Information), where the vesicle is composed
- ⁴⁰ of DOPC:DPPC:Chol:KAON12 = 24:12:24:40 (at this lipid composition, the percentage of the one-phase vesicles with *trans*-KAON12 is 81 % as shown in Fig. 1c). Before UV

irradiation, the *trans*-membrane surface is homogeneous without domains. Immediately after the azobenzene isomer is 45 switched from the trans to cis form by irradiation with UV light, many small domains appear over the entire membrane (Fig. 2a upper). The generated domains exhibit random thermal motion, and the domains become larger through collision and fusion during such motion. Thus, the photo-50 isomerization of KAON12 from the trans- to cis-form induced membrane phase separation. The fluorescence intensities of rho-PE on the membrane surface clearly indicate a difference in lateral organization between the one-phase trans-membrane and two-phase cis-membrane (Fig. 2b). Interestingly, the two-55 phase membrane reverted to one-phase by trans-isomerization (i.e., upon the discontinuation of UV irradiation during treatment with green light) (Fig. 2a lower). Thus, cisisomerization induced stabilized lateral domains, and these domains were again destabilized and disappeared under trans-60 isomerization (Fig. 2c).

Next, we focused on two-phase membranes, i.e., phaseseparated vesicles with *trans*-KAON12. Figure 2d shows typical micrograph of the reversible photo-switching of membrane phase organization (also see a movie and pictures 65 of other giant vesicles in the Supplementary Information), where the vesicle is composed of

- DOPC:DPPC:Chol:KAON12 = 24:12:24:40 (at this lipid composition, the percentage of the two-phase vesicles with *trans*-KAON12 is 19 % as shown in Fig. 1c). Before UV $_{70}$ irradiation, the *trans*-membrane surface is covered by domains. Notably, thermal fluctuation of the domain boundary was occasionally observed with this membrane composition (35 % of the two-phase vesicles exhibited such fluctuating domains). When we irradiated the membrane with UV light to induce $_{75}$ *cis*-isomerization of KAON12, additional small domains
- appeared within the membrane surface (Fig. 2d upper): L_o domains emerged in the L_d phase region, and L_d domains appeared in the L_o phase regions. Furthermore, fluctuation of the domain boundary was suppressed under cis-isomerization, ⁸⁰ which indicates that *cis*-membranes have greater line tension than *trans*-membranes. Upon *trans*-isomerization, the produced domains again disappeared, and phase fluctuation recovered (Fig. 2d lower). Schematic illustrations are shown in Fig. 2e. These reversible changes in membrane phase 85 organization in Fig. 2 were observed more than 10 times. Thus, under UV irradiation, initially uniform one-phase membranes show phase separation into two microscopic coexisting phases, whereas initially two-phase membranes produced additional domains within the membranes. This photo-switching in 90 membrane phases can be observed with a molar fraction of KAON12 of above ~30 %.

Photo-responsiveness of membrane phase organization

The responsiveness of membrane phase organization upon photoisomerization, as shown in Fig. 2, largely depended on ⁹⁵ the composition of ternary lipids (DOPC/DPPC/Chol). We characterized the photo-responsiveness of the membrane system as a function of the DOPC:DPPC ratio with different Chol fractions (Fig. 3). Photo-induced changes in membrane phase separation were most frequently observed, provided that



Fig. 2. Photo-induced reversible organization of lateral domains in one-phase (a-c) and two-phase (d, e) bilayer membranes. (a, d) Fluorescence photomicrographs. The time elapsed is shown on each micrograph. (b) Fluorescence intensities (F.I.) for rho-PE of *trans*- and *cis*-membranes along the dashed line in (a). (c, e) Schematic representation of the molecular switching of the membrane organization. Open circles represent DOPC and KAON12, and filled circles show DPPC and Chol.

membrane ternary compositions were close to the miscibility boundary that separates one- and two-phase states. In the situation close to miscibility, thermal fluctuation of the domain boundary was often observed in the two-phase s membrane surfaces, and all of the fluctuating domains responded as shown in Fig. 2d. In contrast, membranes with compositions far from the miscibility boundary do not respond to photo-isomerization. Photo-isomerization of KAON12 has little effect on membranes with a low Chol 10 concentration of 12%.

Discussion

With regard to thermodynamic phase separation, the stability of bilayer miscibility, such as the emergence of raft domains, is determined by the interfacial energy at the domain edge, i.e., ¹⁵ line tension. When the interfacial energy is as low as the thermal energy, domain edge fluctuation is visible optically, which was previously observed in lipid membranes near miscibility critical points.¹¹ As the temperature is increased

toward the miscibility temperature, edge fluctuation increases,

20 and two-phase membranes transition into a one-phase state without domains. Line tension σ changes depending on the difference between the system temperature T and the miscibility critical temperature T_c as $\sigma \sim (T_c - T)^{\nu}$, where ν is the critical exponent.¹¹ In this study, we observed such change 25 in the edge fluctuation associated with photo-induced phase reorganization, which suggests that the molecular switching of line tension may be the mechanism for photocontrolled domain formation. The change in the molecular conformation of KAON12 into its bulky cis-isomer probably increases the 30 miscibility temperature, consequently leading to the increase in line tension. In other words, miscibility boundary shifts in the isothermal phase diagram under photo-isomerization. It has been reported that the hydrophobic mismatch between the two phases elevates the miscibility temperature.¹² Cis-isomer 35 has a looser packing configuration than trans-isomer at the interface, which possibly enhance such hydrophobic mismatch. Further experiments on the quantitative analysis of the change in line tension are underway.



Fig. 3. Responsiveness of membrane phase organization upon photo-isomerization. (a) Phase diagram of ternary DOPC/DPPC/Chol membranes with 40 % *trans*-KAON12. The open and filled circles correspond to one-phase (homogeneous) membranes without domains and two-phase membranes with domains, respectively. The cross indicates no vesicles. (b-e) Percentage of photo-responsiveness as a function of the DOPC:DPPC ratio with each Chol concentration as shown in part (a): 30% (b), 24% (c), 18% (d), and 12% (e). Responsiveness is defined as the number of vesicles that show photo-induced phase dynamics, which include both initially one-phase and two-phase membranes as shown in Fig. 2 (see Supplementary Information), divided by the total number of observed vesicles. The broken lines show the miscibility boundary.

Yasuhara et al., reported that *cis*-isomerization of an azobenzene-modified cholesterol destabilize lateral domains,⁸ which is opposite to our results. This may be attributed to the difference in partitioning membrane phase between these two 5 molecules: our KAON12 is rich in the L_d region, while their azobenzene-modified cholesterol is preferentially distributed

into the L_o region. When molecular packing in the L_d region becomes much looser under *cis*-isomerization, hydrophobic mismatch between the two phases is enhanced. Conversely, ¹⁰ looser molecular packing in the L_o region suppresses the hydrophobic mismatch. Thus, *cis*-isomelization in the L_d phase tends to increase the miscibility temperature to induce phase-separation into domains, while *cis*-isomerization in the

- L_o phase decreases the miscibility temperature to destabilize to domains. Cell membranes are crowded with transmembrane (TM) proteins that are laterally sorted in L_o (raft) or L_d (nonraft) phase regions according to their preferences for membrane order.⁴ Cells may take advantage of the activation of raft or non-raft TM proteins to effectively control
- ²⁰ membrane domains. Differences in mechanical properties, such as membrane tension, of the model membranes from those of the plasma membranes should be taken into consideration and investigated. Further studies of model and actual plasma membranes are awaited in order to fully ²⁵ understand the molecular mechanism to regulate membrane
- organization.

Recently, using giant vesicle systems, Ayuyan and Cohen reported that photo-induced lipid peroxidation induced uniform vesicles to phase separate.⁹ The mechanism is

³⁰ proposed to be the formation of various amounts of oxidation products within the bilayer. Oxidative conformational changes, such as cleavage of the alkyl chains at the site of a double bond, lead to molecular packing arrangement into membrane domains. However, lipid oxidation is a complex process, and ³⁵ it is hard to extract and quantify the products. There remains unknown parameter concerning the change in molecular conformation. Conversely, our membrane system can provide an efficient tool to examine the effect of the changes in membrane materials on self-assembled mesoscopic structures ⁴⁰ because the photoresponsive molecule can be switched just between *trans*- and *cis*-form reversibly.

We found that membranes that were close to the miscibility boundary showed higher photo-responsiveness, such as 76 % (DOPC:DPPC:Chol = 15:15:30) and 74 % (DOPC:DPPC:Chol ⁴⁵ = 21:15:24). The existence of other vesicles that do not response even in the situation close to miscibility is probably due to a wide variation in the mixed fraction of KAON12 during the vesicle preparation, as previously reported.¹⁰

Here, we examined lateral phase separation of multi-50 component membranes with 40 % KAON12 under photoisomerization. We confirmed that this photo-switching in membrane phases can be observed with a molar fraction of KAON12 of above ~30 %. No apparent difference in the photo-induced phase behavior, as shown in Fig. 2, was noted 55 between these different fractions of KAON12. Notably, at a KAON12 concentration above 50 %, we could not obtain any giant vesicles. The necessary amount of KAON12 (~30 %) for the photo-induced microscopic phase organization is nearly equal to the amount for the previously reported shape $_{60}$ transitions of homogeneous giant vesicles: ${\sim}20$ % and ${\sim}30$ % of KAON12 were required to induce the vesicular transformations of closed vesicles and the formation of membrane pores, respectively.10 Trials on the molecular design of photoresponsive amphiphiles toward the elegant 65 optical control of self-assembled membranes with their smaller amount are awaited.

Conclusions

We succeeded in controlling membrane phase separation of

giant vesicles through the action of a photoresponsive amphiphilic molecule (KAON12). Photo-isomerization induces lateral phase separation in one-phase membranes or produces additional lateral domains in two-phase membranes,

- s in a reversible manner. Our results indicate that, provided that membranes are close to miscibility, molecular perturbations, such as isomerization, can lead to a profound change in laterally organized structures to trigger the formation of raft domains. We discussed the mechanism of this photocontrolled
- ¹⁰ membrane phase separation in terms of the change in the miscibility temperature due to the switching of molecular packing configuration. These findings may contribute to a better understanding of the molecular mechanism for regulating plasma membrane rafts, and may lead to new
- ¹⁵ methods for controlling mesoscopic membrane structures. As the next research target, it may of value to try to apply the methodology on the investigation of the interleaflet interactions of heterogeneous bilayer membranes, by preparing asymmetric bilayers with the photoresponsive ²⁰ amphiphile just on one side.¹³

Experimental

Materials

We designed and synthesized a photosensitive amphiphilic molecule containing azobenzene (KAON12); the details of ²⁵ synthesis have been described previously.⁸ The conformation (*trans* or *cis*) of this molecule can be switched by irradiation with light (Scheme 1). 1,2-Dioleoyl-sn-glycero-3phosphocholine (DOPC), 1,2-dipalmitoyl-*sn*-glycero-3phosphocholine (DPPC) and cholesterol (Chol) were

- $_{30}$ purchased from Avanti Polar Lipids (Alabaster, USA). The fluorescent lipid N-(rhodamine red-X)-1,2-dihexadecanoyl-sn-glycero-3-phosphoethanolamine triethylammonium salt (rho-PE, λ_{ex} = 560 nm, λ_{em} = 580 nm) was obtained from Invitrogen. Deionized water, obtained from a Millipore Milli
- 35 Q purification system, was used to prepare reagents.

Preparation of giant vesicles

Giant vesicles were prepared by the natural swelling method from dry lipid films;⁸ lipid mixtures of KAON12, DOPC, DPPC and Chol dissolved in methanol/chloroform (1:3, v/v)

⁴⁰ along with rho-PE in a glass test tube were dried under vacuum for 3 h to form thin lipid films. The films were then hydrated overnight with deionized water at 37 . The final concentration was 0.2 mM of lipids (DOPC/DPPC/Chol/KAON12) and 1mol% rho-PE.

45 Microscopic observation under photoirradiation conditions

The sample solution (3 μ L) was placed on a glass coverslip, which was covered with another smaller coverslip at a spacing of ca. 0.1 mm. We observed changes in membrane morphology with a fluorescence microscope (IX71, Olympus,

⁵⁰ Japan) equipped with a confocal scanner (CUS-10, Yokogawa, Japan), at room temperature. A 532 nm laser was used to excite rho-PE, and for *trans*-isomerization. *Cis*-isomerization was achieved by UV irradiation through a standard filter set (WU, Olympus; λex=330–385 nm, dichroic mirror 400 nm,

ss λ em=420 nm), with the use of an extra-high-pressure mercury lamp (100 W). The molecular conformation (*trans* or *cis*) of KAON12 was switched by photo-irradiation with green light either with or without UV light (see Supplementary Information). The power of the irradiated light on the ⁶⁰ microscopy stage was 72 µW (green) and 552 µW (UV), which was measured using an optical power measurement system (X-Cite XP750 & XR2100, EXFO, Canada). The images were recorded on a hard-disc drive at 30 frames s⁻¹.

Acknowledgements

⁶⁵ This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas "Soft Matter Physics" and "Bio Manipulation" from the MEXT of Japan, and by the Sunbor Grant from the Suntory Institute for Bioorganic Research.

Notes and references

- ^a School of Materials Science, Japan Advanced Institute of Science and Technology, 1-1 Asahidai, Nomi, Ishikawa 923-1292, Japan, Tel: +81-761-51-1650; E-mail: t-hamada@jaist.ac.jp, takagi@jaist.ac.jp
 ^b Graduate School of Engineering, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan
- 75 † Electronic Supplementary Information (ESI) available: pictures of other giant vesicles, percentage of membrane phase and photoresponse for each membrane composition, and movies of photo-induced membrane phase changes corresponding to Fig. 2a and d. See DOI: 10.1039/b000000x/
- 80 1 D. Lingwood, K. Simons, Science 2010, 327, 46.
- (a) K. Simons, E. Ikonen, *Nature* 1997, 387, 569. (b) D. A. Brown, E. London, J. Biol. Chem. 2000, 275, 17221.
- 3 A. Kusumi, I. Koyama-Honda, K. Suzuki, *Traffic* 2004, 5, 213.
- 4 J. F. Hancock, Nat. Rev. Mol. Cell. Biol. 2006, 7, 456.
- (a) T. Baumgart, S. T. Hess and W. W. Webb, *Nature* 2003, 425, 821.
 (b) M. Staykova, R. Lipowsky and R. Dimova, *Soft Matter* 2008, 4, 2168.
 (c) T. Hamada, Y. Miura, K. Ishii, S. Araki, K. Yoshikawa, M. Vestergaard and M. Takagi, *J. Phys. Chem. B* 2007, 111, 10853.
 (d) L. Bagatolli and P. B. S. Kumar, *Soft Matter* 2009, 5, 3234.
- ⁹⁰ 6 A. T. Hammond, F. A. Heberle, T. Baumgart, D. Holowka, B. Baird and G. W. Feigenson, *Proc. Natl. Acad. Sci. U.S.A.* 2005, **102**, 6320.
- 7 A. P. Liu and D. A. Fletcher, *Biophys. J.* 2006, **91**, 4064.
- 8 K. Yasuhara, Y. Sasaki and J. Kikuchi, *Colloid Polym. Sci.* 2008, **286**, 1675.
- 95 9 (a) A. G. Ayuyan and F. S. Cohen, *Biophys. J.* 2006, 91, 2172. (b) L.
 S. Hirst and J. Yuan, *Liquid Crystals* 2009, 36, 739. (c) N. F.
 Morales-Penningston, J. Wu, E. R. Farkas, S. L. Goh, T. M.
 Konyakhina, J. Y. Zheng, W. W. Webb and G. W. Feigenson, *Biochim. Biophys. Acta* 2010, 1798, 1324.
- 100 10 (a) T. Hamada, Y. T. Sato, T. Nagasaki and K. Yoshikawa, *Langmuir* 2005, 21, 7626. (b) K. Ishii, T. Hamada, M. Hatakeyama, R. Sugimoto, T. Nagasaki and M. Takagi, *ChemBioChem* 2009, 10, 251. (c) T. Hamada, R. Sugimoto. M. C. Vestergaard, T. Nagasaki and M. Takagi, *J. Am. Chem. Soc.* 2010, 132, 10528.
- 105 11 (a) A. R. Honerkamp-Smith, P. Cicuta, M. D. Collins, S. L. Veatch, M. den Nijs, M. Schick and S. L. Keller, *Biophys. J.* 2008, 95, 236.
 (b) S. L. Veatch, P. Cicuta, P. Sengupta, A. Honerkamp-Smith, D. Holowka and B. Baird, *ACS Chem. Biol.* 2008, 3, 287. (c) B. L. Stottrup, A. M. Heussler and T. A. Bibelnieks, *J. Phys. Chem. B* 2007, 111, 11091. (d) C. Esposito, A. Tian, S. Melamed, C. Johnson, S.-Y. Tee and T. Baumgart, *Biophys. J.* 2007, 93, 3169.
 - 12 A. J. Garcia-Sáez, S. Chiantia and P. Schwille, J. Biol. Chem. 2007, 16, 33537.
- 13 (a) M. D. Collins and S. L. Keller, *Proc. Natl. Acad. Sci. U.S.A.*2008, **105**, 124. (b) T. Hamada, Y. Miura, Y. Komatsu, Y. Kishimoto, M. Vestergaard and M. Takagi, *J. Phys. Chem. B* 2008, **112**, 14678.
 (c) S. May, *Soft Matter* 2009, **5**, 3148.